





Ca 3. 26

R 31687

ON THE
TREATMENT OF DISEASES OF THE SKIN,

WITH

*AN ANALYSIS OF ELEVEN THOUSAND
CONSECUTIVE CASES.*



ON
THE TREATMENT OF
DISEASES OF THE SKIN:

WITH
*AN ANALYSIS OF ELEVEN THOUSAND
CONSECUTIVE CASES.*

BY
DR. MCCALL ANDERSON,

*Professor of Practice of Medicine in Anderson's University;
Physician to, and Lecturer on Clinical Medicine in, the Royal Infirmary;
Physician to the Dispensary for Skin Diseases, and to the Cutaneous Wards of
the University Hospital, &c., Glasgow.*

London:
MACMILLAN AND CO.

1872.

LONDON :
R. CLAY, SONS, AND TAYLOR, PRINTERS,
BREAD STREET HILL.

CONTENTS.

PART I.

ANALYSIS OF ELEVEN THOUSAND CONSECUTIVE CASES OF SKIN DISEASE.

CHAPTER I.

Classification.—Statistics of 10,000 consecutive cases of Skin Disease met with in hospital practice, and of 1,000 in private practice :—

A. Functional Affections : Pruritus—Seborrhœa—Comedones—Milia—Hyperidrosis—Ehidrosis cruenta—Ephelis—Melanopathia—Vitiligo—Atrophia cutis—Alopecia—Hirsuties—Fragilitas crinium—Canities *Pp.* 3-25

CHAPTER II.

B. Organic Affections : 1. Parasitic Affections : Tinea favosa—Tinea trichophytina—Tinea versicolor—Tinea decalvans—Scabies—Phtheiriasis. 2. Syphilitic Affections : Secondary and Tertiary Accidents—Hereditary Syphilis, Infantile and Non-Infantile. 3. Strumous Affections : Lupus vulgaris and erythematodes—Scrofuloderma *Pp.* 26-42

CHAPTER III.

Inflammations : Eczema—Erythema — Pityriasis — Prurigo — Ecthyma — Psoriasis — Acne vulgaris — Rosacea — Pemphigus — Pityriasis rubra—Lichen ruber *Pp.* 43-54

CHAPTER IV.

New Formations : Varix — Verruca — Scleroderma — Elephantiasis Arabum and Græcorum—Primary Cancer of Skin—Multiple Fatty Tumours—Hæmorrhages—Purpura—Purpura rheumatica. *Pp.* 55-72

PART II.

THE THERAPEUTICS OF DISEASES OF THE SKIN.

CHAPTER I.

Danger of “driving in” an eruption. A. Local Treatment : General indications for the use of Local Applications—Removal of Crusts, &c.—Indications for the use of Soothing Applications—Baths—Hydropathic Treatment—Shower-bath—Absorbent Powders—Poultices—Soothing Ointments and Lotions . *Pp.* 75-89

CHAPTER II.

Local Treatment (*continued*) : Emphyreumatic oils and their derivatives, Tar, Creosote, Carbolic Acid, &c.—Potash Applications—Pfeuffer’s Treatment—Mercurial Applications . *Pp.* 90-100

CHAPTER III.

Local Treatment (*continued*) : Sulphur—Acetate of Soda—Impermeable Dressings—Vulcanized India-rubber, &c. *Pp.* 101-113

CHAPTER IV.

Local Treatment (*continued*): Caustics—Arsenic—Potassa fusa—Nitrate of Silver—Red Iodide of Mercury—Acid Nitrate of Mercury—Actual Caution—Blisters—Ointment of Cantharides—Iodine—Parasitocides *Pp.* 114-129

CHAPTER V.

B. Constitutional Treatment: Preliminary Remarks—Purgatives, Aperients, and Tonic Aperients—Diuretics—Alkalies—Sedatives and Narcotics—Cod Liver Oil—Iodine and its compounds.
Pp. 130-142

CHAPTER VI.

Constitutional Treatment (*continued*): Tonics, Arsenic—Tar—Carbolic Acid *Pp.* 143-158

CHAPTER VII.

Constitutional Treatment (*continued*): Mercury: subcutaneous injections of—Mercurial Vapour-bath *Pp.* 159-165

CHAPTER VIII.

Constitutional Treatment (*concluded*): Iodide of Potassium—Alkaline Sulphites and Hyposulphites *Pp.* 166-173

ERRATA.

At page 38, line 5 from the top, after the words "it is the only case on record,"
read "with the exception of the one mentioned at page 35."

At page 46, line 6 from the bottom, *instead of* "scabies, prurigo, and urticaria,"
read "scabies, phtheiriasis, and urticaria."

PART I.

ANALYSIS OF ELEVEN THOUSAND CONSECUTIVE CASES OF SKIN DISEASE.

ANALYSIS OF ELEVEN THOUSAND CONSECUTIVE CASES OF SKIN DISEASE.

CHAPTER I.

THE object of the present work is to afford statistics of 10,000 consecutive cases of skin disease met with in hospital practice, and 1,000 consecutive cases met with in private practice, to compare the results, and to allude to a few of the more interesting cases. But it must be borne in mind at the outset that no idea can be formed of the actual frequency of the different forms of skin disease from any statistics, seeing that one is most apt to be consulted with regard to those which are associated with derangement of the general health, which give rise to irritation, or which produce deformity ; still less from the pre-

sent statistics, which of necessity contain an undue proportion of obstinate and rare forms of disease.

It is unnecessary to dwell at any length upon the system on which these diseases have been arranged, especially as their classification is but a modification of that adopted many years ago at the Glasgow Dispensary for Skin Diseases as explained by the late Dr. A. B. Buchanan, in a very able article which he communicated to the *Edinburgh Medical Journal* (January, 1863). The object aimed at in this classification was to render it as useful as possible from a clinical point of view, and hence the most important point was to arrange the various diseases as far as practicable in accordance with their nature and cause. It is most desirable, no doubt, to have a classification in accordance with one principle; but in the present state of our knowledge it is impossible to fulfil the latter indication except at the expense of the former; and accordingly two principles are involved in this clinical classification—namely, the etiological and the pathological.

We divide skin diseases, then, into two great

classes—namely : A, Functional ; and B, Organic. The Organic we subdivide into two great classes—I. Those defined by uniform causes ; II. Those not defined by uniform causes. The diseases defined by uniform causes are arranged under four heads—namely : 1, Parasitic affections ; 2, Syphilitic affections ; 3, Strumous affections ; 4, Eruptive fevers. The diseases not defined by uniform causes comprise all affections of the skin not included in any of the preceding groups, and are arranged pathologically under three heads—namely : 1, Inflammations ; 2, New formations ; 3, Hæmorrhages.

CLASSIFICATION OF 10,000 CONSECUTIVE CASES OF SKIN DISEASE MET WITH IN HOSPITAL PRACTICE.

A.—FUNCTIONAL AFFECTIONS.

Pruritus	39
Seborrhœa	13
Comedones	10
Milium	1
Hyperidrosis	5
Ephelis	3
Melanopathia	1
Vitiligo	4
Atrophia cutis	2
Alopecia	56
Hirsuties	2
Fragilitas crinium	3

B.—ORGANIC AFFECTIONS.

I. DISEASES DEFINED BY UNIFORM CAUSES. |

1. *Parasitic Affections.*

Vegetable	{	Tinea favosa	156
		{	Circinata 54
			Tonsurans 67
			Circinata + tonsurans . . . 3
			Sycosis 18
		Tinea versicolor	106
		Tinea decalvans	153
Animal	{	Scabies	2527
		Phtheiriasis	327

2. *Syphilitic Affections.*

Primary accidents	24
Secondary and tertiary accidents	437
Hereditary syphilis (infantile)	55
„ „ (non-infantile)	1

3. *Strumous Affections.*

Lupus 198	Strumous glands 191
Scrofuloderma 27	„ ulcers 85
Lichen scrofulosorum 3	„ abscess 8

4. *Eruptive Fevers.*

Morbilli	1
Scarlatina	2
Variola	6

II. DISEASES NOT DEFINED BY UNIFORM CAUSES.

I. *Inflammations.*

Erythema ¹	470	Erysipelas	10
„ multiforme	1	Acne vulgaris	288
„ nodosum	5	Rosacea	37
„ pernio	2	Psoriasis ⁵	725
Eczema ²	2527	Pemphigus	12
Impetigo contagiosa	10	„ foliaceus	1
Ecthyma ³	97	Pityriasis rubra	6
Prurigo	1	Deep inflammations	345
Urticaria ⁴	147	Ulcers ⁶	433
Zona	32	Onychia	10
Dermatitis	27		

2. *New Formations.*

Varix (uncomplicated)	41	Elephantiasis Græcorum	1
Nævus vascularis	9	Molluscum contagiosum	6
„ spilus	2	Fibroma molluscum	1
Verruca	29	Cicatrix	4
Callositas	10	Cheloid	5
Clavus	1	Cysts	23
Cornu	2	Scirrhus	5
Ichthyosis	31	Epithelioma	38
Scleroderma	1	Unclassified tumours	2
Elephantiasis Arabum	7		

3. *Hæmorrhages.*

Purpura simplex	6
„ tuberculosa	1
„ rheumatica	1

¹ Including strophulus, pityriasis, and roseola.² Including its lichenous, pruriginous, and impetiginous forms.³ Including rupia non-syphilitica.⁴ Including lichen urticatus.⁵ Including lepra.⁶ Independent of struma and syphilis.

CLASSIFICATION OF 1,000 CONSECUTIVE CASES OF SKIN DISEASE MET WITH IN PRIVATE PRACTICE.

A.—FUNCTIONAL AFFECTIONS.

Pruritus	11	Atrophia cutis	1
Comedones	7	Alopecia	18
Hyperidrosis	1	Hirsuties	5
Ephidrosis cruenta	1	Fragilitas crinium	2
Ephelis	3	Canities	4

B.—ORGANIC AFFECTIONS.

I. DISEASES DEFINED BY UNIFORM CAUSES.

1. *Parasitic Affections.*

Vegetable	Tinea favosa epidermidis			4
	Tinea trichophytina	Circinata ¹	10	36
		Tonsurans	20	
		Sycosis	6	
Animal	Tinea versicolor			15
	Tinea decalvans			44
	Scabies			44
	Phtheiriasis			7

2. *Syphilitic Affections.*

Secondary and tertiary accidents	51
Hereditary syphilis (infantile)	3
„ „ (non-infantile)	3

3. *Strumous Affections.*

Lupus	{ Vulgaris 22 Erythema- todes 3 }	. 25	Lichen scrofulosorum	2
			Strumous glands	9
			„ ulcers	2
Scrofuloderma		6	„ abscess	1

¹ Including the so-called eczema marginatum.

II. DISEASES NOT DEFINED BY UNIFORM CAUSES.

1. *Inflammations.*

Erythema ¹	99	Rosacea	21
„ pernio	2	Psoriasis ³	106
Eczema ²	348	Pemphigus	3
Impetigo contagiosa	1	Pityriasis rubra	1
Prurigo	2	Lichen ruber	1
Urticaria	5	Furunculus	10
Zona	1	Ulcers ⁴	7
Acne vulgaris	54		

2. *New Formations.*

Varix (uncomplicated)	4	Elephantiasis Arabum	1
Nævus vascularis	2	„ Græcorum	1
„ spilus	1	Scirrhus	2
Verruca	4	Epithelioma	9
Ichthyosis	7	Multiple fatty tumours	1
Scleroderma	1		

3. *Hæmorrhages.*

Purpura rheumatica	1
------------------------------	---

A.—FUNCTIONAL AFFECTIONS.

The *functional* affections comprise all those diseases of the skin and its appendages in which

¹ Including strophulus, pityriasis, and roseola.

² Including its lichenous, pruriginous, and impetiginous forms.

³ Including lepra.

⁴ Independent of struma and syphilis.

there is no evidence of structural change. Of these there were 192 cases, 53 of which occurred amongst the private, and 139 amongst the hospital cases, and they include fifteen separate affections.

The first of these, pruritus, of which there were 50 illustrations, includes only those cases in which itching was the sole cutaneous symptom, and which may arise from a variety of causes, such as disorders of the digestive organs, of the urinary and genital organs, and of the nervous system; and the treatment consisted in the removal of the cause or causes, in addition, frequently, to the empirical treatment of the symptom itself. There can be little doubt that very frequently cases are erroneously classed under this head: thus the pruritus on careful investigation is often found to be dependent on the ravages of the itch insect or the louse, or upon recurrent outbreaks of nettle-rash, and should then be classed under the respective heads of scabies, phtheiriasis, and urticaria. The only case specially worthy of mention under this head is that of a lady, aged about fifty, who

consulted me on Jan. 10th, 1870, on account of intolerable irritation of the skin, which deprived her in great measure of sleep, and made "her life a burden to her." The history which she gave of her case was that she went to visit a friend who was very deaf, and who used a speaking trumpet. She put her mouth close to the mouth-piece of the trumpet, and immediately experienced the irritation complained of, and from that time to this, a space of nearly two years, it has never left her. No trace of disease can be detected in her skin, and the pruritus cannot be very severe, as there are no marks of her nails to be seen. Her general health appears to be good, but she is in a very depressed state, and fears that she will never be better. It is quite evident that in this case the disease is mental rather than physical, and it is not improbable that it may end in insanity.

Of the functional affections of the sebaceous glands there were 31 illustrations, representing three forms of derangement—namely, seborrhœa, comedones, and milium. Under the head of seborrhœa, of which there were 13 instances, are

classed those cases in which the sebaceous glands were abnormally active, and in which, as a consequence, an excessive quantity of sebaceous matter appeared on the surface in the fluid form or in the shape of oily crusts.

The term "comedones," of which there were 17 examples, refers to accumulations of hardened sebaceous matter in the ducts of the sebaceous glands, or in the hair-follicles communicating with them, their presence being indicated by little black specks upon the skin, due to admixture of the sebaceous matter on the surface with particles of dust. Cases in which these accumulations became complicated with inflammation of the skin in their vicinity have been classed under the head of acne.

Milium, for which only one patient came specially for advice, but which was met with incidentally many times in patients complaining of other forms of skin disease, differs from comedones in this respect, that, owing to an obstruction in the duct of the gland, the latter, in whole or in part, becomes distended with seba-

ceous matter, thus forming a little pearly-looking elevated spot upon the surface.

Functional affections of the sudoriparous glands were more rarely met with, there having been only six cases, all of which were examples of hyperidrosis or excessive perspiration, and in one at least of these the perspiration was accompanied by fœtor. The following extract from a note received from a surgeon in England affords a good instance of this disagreeable affection:—"I suffer from axillary perspiration to a very disagreeable extent; so much so that often when I am walking about seeing patients my shirt and coat in the region of the axilla are perfectly wet. This occurs even on the coldest day. My skin in other parts is remarkably free from perspiration. I need hardly say that I observe the most perfect cleanliness."

It may be well in this place to refer to that extraordinary complaint improperly termed ephidrosis cruenta, or bloody sweat, for it is not a perspiration at all, but an exhalation of blood from the pores of the skin. The rarity of this affection is proved by the fact that it only occurred

once out of the whole 11,000 cases. It is unnecessary to give more than a mere outline of this case, as it, along with similar cases, was fully described by the writer in the *Journal of Cutaneous Medicine* (vol. i. page 328). The patient, a young lady fourteen years of age, was seen by me at the request of Dr. Mason, of Ayr, in the summer of 1866. The parts implicated were the face, arms, front of the chest, and legs. The hæmorrhage occurred from round erythematous patches of eruption, which were remarkable for their symmetry. One was on the brow, another on the chin, and one on each cheek. On the front of each arm also there were four in a row, two on each upper arm and two on each fore-arm. A similar arrangement was observed upon the sternum and upon the legs. One of the most marked peculiarities of the hæmorrhage was the suddenness of its invasion. An oval or round red ring, varying from the size of a shilling to that of a crown, formed almost instantaneously, and the redness quickly spread inwards over the enclosed skin. As soon as seen the patches appeared as if

the cuticle had melted away, and the surface was quite wet. Sometimes the exudation was like water at first, and changed into blood; at other times, and especially on the face, the patches were at once covered with a complete dew of blood. The hæmorrhage did not, however, consist merely of the dew of blood; that was only at the outset: it was actual bleeding, as from a cut, the blood sometimes streaming down the face or other part attacked. There was rarely more than one attack each day, although sometimes the bleeding occurred from two separate portions of skin simultaneously. It is very curious to note that the outbreak *generally* occurred at the same hour each day—namely, at 11 A.M.; but it did not seem to be under the influence of mental or bodily excitement, or to be induced by taking food or stimulants. It was evident that the hæmorrhage was dependent upon defective and irregular menstruation—that, in fact, it was a case of vicarious menstruation; and accordingly it yielded, in the space of three or four weeks, to remedies directed against the latter disorder.

Of pigmentary disorders of the skin there were 11 cases, including three different forms of disease—namely, ephelis, melanopathia, and vitiligo. As these terms have been variously used by dermatologists, it may be right to explain the meaning that is attached to each in the present classification.

By ephelis is meant an excessive deposit of pigment in the mucous layer of the epidermis, giving rise to brown patches of various shapes and sizes, such as we so frequently observe upon the brow of the pregnant female. Of this there were six cases, three amongst the 1,000 private, and three amongst the 10,000 hospital cases.

There was only one case classed under the head of melanopathia, by which is meant an irregular and streaky deposit of pigment giving to the surface a marbled appearance. This condition is most frequently met with upon the legs, as the result of long-continued exposure to the heat of the fire ; but it occurs in its most typical form in persons tainted with syphilis. Such cases, however, of which a good many presented themselves, are classed under the head of constitutional syphilis.

In vitiligo, of which four cases came under observation, there is not so much an excessive deposit as an irregular distribution of the pigment, so that variously shaped, unduly white patches, at the edges of which the skin is unnaturally dark, are presented to view.

It may be well at this time to refer to a case which, as far as known to the writer, is unique, in which the symptoms were at first identical with those of *tinea decalvans* (*alopecia areata*), but subsequently with vitiligo. A young lady, aged ten, healthy-looking, and born of a sound stock, consulted me on June 30th, 1870, on account of round and irregular bald patches on the head, the latter being due to the coalescence of neighbouring round ones, and implicating, in all, about one-half of the head. The case, in fact, presented all the naked-eye characters of *tinea decalvans*, the disease due to the presence of the *microsporon Audouini*; but, unfortunately, no note was taken at the time as to the result of the microscopic examination of the hairs. She was recommended to regulate the bowels with simple

aperients, to take small doses of wine of iron and Fowler's solution, and, after shaving the head, to sponge it night and morning with a lotion of perchloride of mercury (four grains to the ounce). On July 28th all the bald patches were thickly clothed with hair, which, as is usual in cases of tinea decalvans when the hair first reappears, was white, owing to the absence of pigment. So far there was nothing unusual in the symptoms until Dec. 23rd, when the patient again visited me. The hair was then perfectly healthy, but as white in the sites of the previous bald patches as on the 28th of July, the scalp in these situations being likewise devoid of pigment. She then showed me what had appeared, about a fortnight previous, on her shoulders and back—namely, round and oval white spots, from the size of a crown downwards, the skin at the edges of which was deeply pigmented. In fact, she now presented all the characteristic symptoms of vitiligo.

Vitiligo is in most cases either a congenital or nearly congenital deformity; but there are exceptions to this rule, as in the following case:—

Mary M——, aged twenty-seven, domestic servant, came to the Glasgow Dispensary for Skin Diseases on the 27th January, 1869, on account of a discoloration of the skin. Without questioning, she volunteered the following statement as to the origin of the disease:—"About eight years ago, whilst working in a hay-field, I was struck on the back of the head by a fellow-worker so forcibly as to drive a hair-pin into my head." About a year after this her attention was called by a companion to a patch of grey hair on the back of her head; and, on close examination, the scalp in this situation was found to be white in colour. Shortly after this she noticed many white spots on the sides of her neck. On examination of the patient, a swarthy-looking Irishwoman, the following appearances were noticed:—On the occipital region and upper portion of the neck, in the median line, the hair, to the extent of nearly four inches in length and two in breadth, was quite grey, and the skin from which it grew of a clear white tint; while at the edges of the patch, which were irregular in form, the skin was

much darker than the healthy skin in the vicinity. On the sides of the neck above the clavicles were two patches four inches long and one broad, presenting exactly the same characters minus the hair. Similar patches, varying in size from a threepenny-piece to a florin, were seen on the chest and right arm. Her general health appeared to be excellent. The formation of new patches, which was going on at first, was arrested, and some improvement in the existing ones was observed, under the influence of Fowler's solution in small doses continued for some months.

Of simple atrophy of the skin there were three cases, two of them occurring in hospital and one in private practice. All of them were very circumscribed. In two the atrophy consisted of a narrow, white, depressed line, extending from the supraorbital notch nearly perpendicularly over the brow along the track of one of the branches of the supraorbital nerve. In the third case the atrophy implicated small irregular patches of skin upon the leg, brow, upper lip, and neck below the right ear. This is evidently a neurotic affec-

tion, but in none of the cases alluded to could any satisfactory explanation be given of its occurrence.

Among the 90 cases of functional disorder of the hair or hair-follicles, 29 of which occurred among the private patients, there were four different forms of disease—namely, alopecia, hirsuties, fragilitas crinium, and canities.

The term “alopecia,” of which there were 74 examples, is used in this classification, as it should always be, in a very restricted sense. Thus, when the scalp is attacked by erythema, the hair often comes away in great abundance; but here the alopecia is merely one of the symptoms of erythema (or pityriasis, as it is generally termed). When the poison of syphilis enters the blood, the hair frequently falls out; but this is merely one of the manifestations of syphilis. When lupus attacks the head, it destroys the hair-follicles and leads to permanent baldness; but the alopecia in that case is merely one of the consequences of the strumous disease. And, lastly, when the microsporon Audouini attacks the hair, it produces bald

patches; but this is only one of the symptoms of tinea decalvans. Such cases are classed under the respective heads of erythema, syphilis, lupus, and tinea decalvans. Idiopathic alopecia usually occurs as the result of debility—during convalescence from a fever, for example; although it is sometimes hereditary; and occasionally, as I shall endeavour to point out when referring to the pathology of tinea decalvans, a purely neurotic affection.

There were seven cases of hirsuties, five of which occurred among the private cases. By this is meant an excessive development of hair in unusual situations, especially in females. There is much difference of opinion as to the proper advice to be given to such persons. Some recommend the extraction of the hairs, under the belief—which is shared in by many members of the profession, even the most distinguished—that when a hair is extracted along with its bulb, it does not grow again. But this is the worst recommendation of all; for not only does it grow again, but the stimulation of the epilation makes it grow with

greater vigour than before. The treatment of hirsuties therefore resolves itself into the regular employment of the razor or of a depilatory. The latter is certainly to be preferred, provided the agent used is carefully selected. In doing so, we should aim at securing an application which is efficient without being irritating ; and the remedy which I have found to fulfil these conditions in a satisfactory manner is composed of—sulphuret of barium, a drachm and a half ; oxide of zinc, six drachms ; carmine, one grain : this is mixed with a sufficient quantity of water to form a paste, which is smeared over the hairy part ; in three minutes it is washed off, and the hair comes away with it.

The next disease of the hair, which has been named “*fragilitas crinium*” by Wilson, and of which five cases came under observation, is little known to the profession. The whiskers and moustache are more frequently affected than the hair of the head. The affected hair is marked by a variable number of white spots, and after it grows a certain length it has a tendency to break at

one of these points. On microscopic examination it is found that the white spots are partial fractures of the hair, their nutrition being evidently defective, and their elasticity proportionately impaired.

Of canities, or decoloration of the hair, there were four instances, all of which occurred amongst the 1,000 private cases. One of these is specially worthy of note. This was the case of a gentleman in the prime of life, and apparently in good health. The affection was of some years' duration, and implicated the hair of the left upper eyelid and most of the hair of the left whisker, which was quite white. The affection of the eyelid was complete within a few days from its commencement, the whiteness appearing first at the inner angle of the lid, and rapidly extending outwards. No cause could be assigned by the patient, unless anxiety in starting a new business. In some cases the decoloration is hereditary; thus I am acquainted with a gentleman who has jet-black hair with the exception of one lock of grey in the occipital region, who informed me

that his father and grandfather presented the same peculiarity. In other cases the whiteness of the hair is merely to be regarded in the light of premature disappearance of the pigment, such as occurs in the natural course of events in aged persons.

CHAPTER II.

B.—ORGANIC AFFECTIONS.

OF the *diseases defined by uniform causes*, the parasitic affections, which constitute the first class, bulk largely in the present statistics; for of the 11,000 cases, no less than 3,561 were dependent upon the presence of parasites. These affections have been divided, as usual, into two sub-classes—namely, (1) affections due to the presence of vegetable, and (2) affections due to the presence of animal, parasites. With regard to the first of these, the vegetable parasitic affections, which were 656 in number, there is much difference of opinion, and consequently much variety in their classification. Thus, Wilson holds that there are no such diseases, the plant-

like structures met with in ringworm, &c., being, in his opinion, mere degenerations of the structures of which the skin is composed. Others, while admitting the class of vegetable parasitic affections, hold that they are all due to one and the same parasite; while my own opinion, as fully explained elsewhere, and as indicated in the present statistics, is that, regarded from a clinical point of view, there are four fungous growths, productive of four distinct affections, which under no circumstances are transmutable.

The first of these, *tinea favosa*, due to the presence of the *achorion Schönleini*, seems to be much more frequent in Scotland than in England, having been met with no less than 160 times. It rarely occurs except amongst the lowest classes, as is proved by the fact that no less than 156 of the cases occurred in hospital practice, while the four which were met with in private were all in members of the same family, amongst whom it spread by contagion. In this, and in many other instances, the disease was contracted from the lower animals

—in most of them directly or indirectly from mice. The frequency of favus in mice, coupled with the mousy odour which is one of the characteristics of the complaint, renders it not improbable that favus was originally communicated to human beings from them.

The next vegetable parasitic affection, ringworm, or *tinea trichophytina* (due to the presence of the *trichophyton*), was met with 178 times—36 times amongst the 1,000 private, and 142 times amongst the 10,000 hospital cases. These statistics would lead one to the belief that, in Scotland at least, *tinea favosa* is nearly as common an affection as ringworm, than which nothing could be more erroneous, for there is little doubt that ringworm occurs at least twenty times more frequently than favus; but patients are not nearly so likely to seek advice with regard to the former affection, partly because it is less obstinate and less disfiguring, and partly because of the numerous popular remedies which are in vogue for its removal. It will be observed that, in the opinion of the writer,

tinea circinata, tinea tonsurans, and tinea sycosis are mere varieties of ringworm, the first affecting non-hairy parts, the second the head, and the third the other hairy parts of the body, especially the beard. It may be right, too, to mention that under the head "tinea circinata" are classed some cases of that disease first described by Hebra as a variety of eczema, under the name of eczema marginatum, it being a mere variety of ringworm of the body, as I have endeavoured to prove in a recent number of the *Edinburgh Medical Journal* (May, 1868).

The third vegetable parasitic affection, tinea versicolor, that brown patchy eruption which occurs upon the trunk of the body, due to the presence of the microsporon furfur, was met with 121 times, and with almost equal frequency in all classes of the community; but these statistics afford no criterion of its actual frequency, for patients comparatively rarely seek advice with regard to it. There were numerous illustrations of its contagious nature, and many cases also in which it did not seem to be com-

municable, for all persons are not equally liable to be infected.

The last of the vegetable parasitic affections, *tinea decalvans* (due to the presence of the micro-*sporon Audouini*), occurred 197 times, 44 of which cases were met with among the 1,000 private cases. These statistics would lead one to suppose that it is a commoner affection than *tinea versicolor*: but in reality it is a comparatively rare disease; one, however, which, by destroying the hair and thus producing much deformity, creates alarm, and hence most cases come under the notice of the physician. There is much difference of opinion as to the pathology of this complaint, but I have no doubt of its parasitic nature; and the reason why the fungus is so generally overlooked is that it is very small compared with the other vegetable parasites, and that it implicates the sheath of the hair rather than the hair itself, so that when the latter is extracted without the sheath, as so often happens, no fungus is discovered. At the same time it comes to be a question whether a few of the cases presenting

the naked-eye characters of *tinea decalvans* are not rather the result of perverted innervation than of the presence of a fungous growth. The case referred to under the head of vitiligo (see page 17) lends support to this view.

It is quite evident that all persons are not equally under the influence of vegetable parasites, a certain soil, as in the case of the higher plants, being necessary in order to their growth; and, as illustrated by the cases under review, debilitated and strumous persons seem specially liable to be attacked. Hence constitutional treatment, with the view of improving the general health, was in most cases resorted to, in combination with local measures. It is not improbable that in some of these cases the constitutional treatment, by rendering the soil unsuitable, might have served to remove the disease, but it was not considered desirable to trust to it alone. The local treatment, which in some instances was resorted to exclusively, consisted in the use of parasitocides, such as the perchloride of mercury (two to four grains to the ounce), the

hyposulphite of soda (half a drachm to a drachm, to the ounce) and sulphurous acid. Sometimes the eruptions—erythematous, eczematous, &c.—produced by the irritation of the fungus, had a tendency to remain after the parasiticides had completed their work of destruction, and then the subsequent treatment of these eruptions was conducted on the same principles as if they had occurred independent of such special causes. Where hairy parts were involved it was generally found necessary to extract the hair in addition to using parasiticides, with the view of removing, along with the roots, the fungous matter which lay embedded in the follicles and beyond the reach of the parasiticides. This was especially necessary in the case of *tinea favosa* and *tinea sycosis*. In obstinate cases of *tinea tonsurans* and *tinea decalvans* strong local applications—including blistering—were employed successfully.

The parasitic affections due to the presence of animal parasites were 2,905 in number, and included two forms of disease—scabies and phthieriasis. Of the first, 2,527 occurred amongst the

10,000 public, and 44 amongst the 1,000 private cases ; so that, while scabies is by no means so rare amongst the upper classes as many suppose, occurring to the extent of fully 4 per cent. of the cases under observation, it is vastly more frequent amongst the lower, occurring to the extent of fully 25 per cent.

Of the three varieties of phtheiriasis—*P. capitis*, that which affects the head ; *P. pubis*, that which affects all hairy parts except the head, but especially the hair of the pubes ; and *P. corporis*, that which affects the non-hairy parts ; and each due to a separate variety of *pediculus*—that which implicates the body was by far the most frequently observed, but the exact proportion cannot be stated, as the cases were not kept separate in the present statistics. This disease occurred almost exclusively amongst the lower classes, 327 cases having occurred amongst them, and only 7 amongst the private cases ; and it was generally found, as in the case of the vegetable parasitic affections, that constitutional treatment, such as cod-liver oil and tonics, in addition to the

use of parasiticides, such as ointment of staphisagria and lotions of perchloride of mercury, accelerated the cure and rendered it more permanent.

The proportion of syphilitic cases indicated by the present statistics is somewhat less than had been anticipated, there having been 574 in all, or nearly 5 per cent. ; of which 517 occurred amongst the 10,000 hospital, and 57 amongst the 1,000 private patients. It was not thought necessary to separate the various forms of secondary and tertiary eruption (488 in number) : partly because these are named, somewhat arbitrarily, from their supposed resemblance to non-syphilitic affections, and have, moreover, a tendency to run into one another ; and partly because the treatment adopted depended not so much upon the mere character of the eruption as upon the constitution of the patient and the interval that had elapsed since the contraction of the disease. While, however, mercurial treatment was generally resorted to in cases of secondary, and iodide of potassium in cases of tertiary syphilis, the former was frequently employed

with success in obstinate cases of tertiary syphilis after the latter had failed. Simple tonic treatment was often made use of when the state of the general health appeared to demand it, but it was not relied upon alone in the general run of cases; and, indeed, the results of the treatment, of which the above is an outline, are altogether opposed to the too fashionable belief that anti-syphilitic treatment is unavailing, and that tonic treatment and a reliance on the "*vis medicatrix naturæ*" is all that can be desired.

Of hereditary syphilis there were 62 examples, almost all of which occurred in the infantile period. Of the non-infantile cases, of which hardly any notice is taken by authors, one or two may be alluded to. One of these was a case of pemphigus occurring in a little girl about nine years of age. The eruption of bullæ was somewhat circumscribed, being most abundant about the privates, around the mouth, and on the arms and legs. The eruption assumed the form of segments of circles, as syphilitic eruptions so often

do ; the stains left by it were somewhat coppery ; and there was a distinct history of miscarriages in the mother. She improved greatly under a course of grey powder.

A little boy, aged nine, was brought to me on Feb. 26th, 1858, labouring under an eczematous eruption which had continued since he was six weeks old. The eruption was situated principally upon the face, but a few scattered patches were observed upon the legs. It had all the usual characters of simple chronic eczema : the skin was considerably thickened and infiltrated, and the patches were red, very itchy, and exuded clear serum from time to time. His face was pale, and he had chronic bronchitis ; but the digestive organs were healthy. I saw him from time to time, and treated him at first with tar capsules on account of the bronchitic complication ; subsequently with iron, arsenic, and cod-liver oil ; while the external treatment consisted of various preparations of tar and of mercury in the shape of lotions and ointments. The eruption improved very much, but never disappeared. He was

brought to me again on Oct. 14th, when, disappointed at the failure of what may be termed the anti-eczematous treatment, I examined him carefully. I found that his face was pale and was the seat of small scars; his nose, too, was somewhat depressed, and slight fissures, which were of old standing, were detected at the angles of his mouth. I came to the conclusion that the eruption was the result of an hereditary syphilitic taint, and gave him a teaspoonful three times a day of three drachms of iodide of potassium dissolved in six ounces of infusion of quassia. This did no good, and accordingly on March 3rd, 1869, I prescribed five grains of grey powder twice daily. On the 17th the eruption was greatly better, and the powders had agreed with him. On the 12th of April it had completely disappeared, with the exception of slight redness and roughness of the upper lip, while the fissures at the angles of the mouth were healed, whitish marks only being left. The grey powder was stopped, and I have not seen the patient since, so that, judging from the anxiety of his mother

as to his complaint, it is reasonable to infer that he continues well.

In this case the iodide of potassium failed ; but it is not always so, as in the following case, which is interesting, not only on this account, but also because it is the only case on record of hereditary syphilitic pemphigus (of course, excepting infantile pemphigus, of which many cases have been published).

Master H——, aged about ten years, was brought to me on Oct. 26th, 1867, on account of an eruption of bullæ. These commenced to form on Sept. 14th, and the complaint was perpetuated by the occurrence of fresh crops. At the date of his visit two bullæ were detected, and the remains of one or two others on the hands and feet. No other parts of the body were ever attacked. He was pale, thin, and weakly, and his bowels were too loose. His father died at the age of twenty-eight years, said to be of inflammation of the chest ; and his mother, immediately after his birth, became paralytic, and died at the age of twenty. Taking into consideration all the

facts of the case, and especially the pallor of the countenance and the limitation of the disease to the hands and feet, which I have observed to be the favourite seats of the eruption in syphilitic pemphigus when it is not hereditary, I prescribed—iodide of potash, half an ounce; ammonio-citrate of iron, two drachms; syrup of ginger, two drachms; infusion of cascarrilla, to twelve ounces: a dessert-spoonful three times a day in a wine-glassful of cold water. He was ordered a good deal of animal food, and to abstain from porridge, cheese, pastry, and trash of all kinds.

Immediately after his visit to me he was sent to school in Edinburgh, where the medical attendant of the boys saw him, and was shown my prescription. He evidently took a different view of the case, and therefore very naturally disapproved of the treatment proposed, and gave him instead a course of cod-liver oil. On his return from school for the Christmas holidays he was not the least improved in any respect. His guardian therefore stopped the oil and began

the treatment which I had suggested. After he had finished one bottle of the medicine he was brought to me. He was still pale, but felt very much stronger, and the eruption had quite disappeared, only one or two discoloured marks indicating the seats of the previous bullæ. He was ordered to continue the treatment, and I have not seen him since.

Strumous affections, of which there were 557 examples, are nearly as numerous as syphilitic, if we include strumous glands, strumous abscesses, and strumous ulcers; but if we exclude them, it will be found that they are much less numerous, there being only 261 amongst the whole 11,000 cases. An attentive observation of the cases of lupus afforded no corroboration of the view that that disease is not essentially a strumous one, as many assert; and the fact that sometimes no other manifestation of the strumous diathesis was present is no better argument against its strumous character than is the absence of any other strumous manifestation against the strumous character of strumous adenitis. Of the two

varieties of lupus, lupus erythematodes and lupus vulgaris, the former was only exceptionally met with; while lichen scrofulosorum, the papular eruption described by Hebra, seems to be rarer in Scotland than in Austria, if we may judge by the present statistics, only five undoubted cases having been noted. Strumous affections not belonging to lupus or lichen have been classed under the head of "scrofuloderma," of which there were 33 cases, and in which is included that variety described by me under the head of "scrofuloderma verrucosum."¹ The constitutional treatment of this group consisted in the use of the usual anti-strumous remedies, such as cod-liver oil, steel, sea-air, and attention to the general health; while, locally, caustics were for the most part relied upon, of which the solid nitrate of silver and the acid nitrate of mercury were most frequently used. The constitutional treatment was found most useful in preventing relapses; the local, in removing existing manifestations.

It is unnecessary to dwell upon the last group

¹ Journal of Cutaneous Medicine, vol. i. p. 26.

of diseases defined by uniform causes, the eruptive fevers ; as these, for obvious reasons, were only exceptionally observed at the dispensary, while those occurring in private practice were excluded from the present statistics.

CHAPTER III.

THE first class of diseases not defined by uniform causes—the Inflammations—is a very extensive one, embracing as it does more than one-half of all the cases (namely, 5,847), and this notwithstanding the exclusion of all inflammatory affections included under previous groups on account of their dependence on uniform causes. It is impossible to do more than refer briefly to a few of the more important of these.

The most frequent by far was Eczema, of which there were 2,875 cases (2,527 having occurred amongst the 10,000 hospital, and 348 amongst the 1,000 private patients); that is, in the proportion of rather more than 1 to 4. This is a proportion lower than is usually recognized (namely about one-third), although the term “eczema”

was used in its widest sense, including the so-called lichen and impetigo, the reasons for which have been given at length elsewhere. The proportion would probably have been somewhat greater had it not been that under the head of "scabies" was included, not only those patients who came themselves for advice, but also other members of their families who were reported to be similarly affected. It would be quite out of place to enter into details with regard to this protean affection ; but it may be remarked that of the two forms of eczema, the generalized and the localized, the latter was the more frequently observed, while the former was more readily amenable to treatment. No case of universal eczema—by which is meant an eruption covering the whole of the body without leaving a single interval of sound skin—was observed. Of the localized forms, the parts most frequently attacked were the legs, hands, head, and hairy portions of the face ; and next to them the genital organs, anus, ears, arms and neck, and internal aspect of thighs. In most cases debility, or digestive derangement, or local irritation, or a

combination of these causes, lay at the root of the disease.

Erythema, the simplest of all the inflammatory affections of the skin, was much less frequently met with (namely, 470 times amongst the hospital, and 99 times amongst the private cases), although under it were included cases of the so-called strophulus, non-syphilitic roseola, and pityriasis. A good many cases, however, were classed under eczema which might have been ranged under erythema, as it is sometimes difficult to say where erythema ends and where eczema begins ; for most of the forms of erythema are but lower grades of inflammation than eczema—indeed, most cases of the latter commence, and almost all of them terminate, in erythema.

In the present classification no such disease as pityriasis is recognized ; for if we exclude seborrhœa sicca, sometimes termed pityriasis, and the genuine pityriasis rubra as first described by Devergie, the remaining cases of the so-called pityriasis are nothing more nor less than the second or scaly stage of erythema ; and it is manifestly

absurd to speak of the second stage of a complaint as a separate disease, though it may be convenient to retain the term as applicable to the chronic stage of simple inflammation of the skin.

The disease so graphically described by Hebra under the name of prurigo was very rarely met with—much more rarely apparently than at Vienna, if we may judge from the statistics of the great German dermatologist—perhaps because the lower classes in this country live under better hygienic conditions; but pruriginous eruptions were very commonly observed. By a pruriginous eruption, I mean an eruption produced by the nails of the patient in scratching, such as we notice so generally in cases of scabies, phtheiriasis, and urticaria. Such eruptions are mere symptoms or complications of these diseases, and are therefore classed under their respective heads of scabies, prurigo, and urticaria; while most of the others, which appeared to be independent eruptions, were grouped with eczema (*E. pruriginosum*), always excepting what we may call the true prurigo of Hebra, which, as just observed, was very rarely met with.

Under the head of Ecthyma 97 cases are noted as having occurred amongst the hospital patients, whilst not a single case was observed amongst the private ones. We are told by some dermatologists—Hebra for example—that there is no such disease as ecthyma, the cases previously classed under this head being mere symptoms or complications of other diseases, such as syphilis, scabies, and eczema. But while it must be admitted that many cases are described as ecthyma which in reality have no title to such a classification, the present statistics do not confirm the view just mentioned. Under the head of ecthyma have been classed only those cases in which large isolated pustules terminating in crusts and occurring generally in debilitated subjects were observed, independent of syphilis, of scabies, and of all other itchy affections. The constitutional treatment, consisting of generous diet, change of air, and the use of tonics, was most relied upon, while soothing local applications gave much relief and accelerated recovery.

The most frequent of all the inflammatory affections, with the exception of eczema, was

Psoriasis, of which there were 831 examples, 725 of which occurred amongst the hospital cases, and 106 amongst the private ones. This is a much larger number than had been anticipated at the outset, but which is perhaps accounted for in part by the fact that psoriasis is not only a very disfiguring and very obstinate affection, but also one which has a marvellous tendency to recur, especially in the spring and autumn months. It must be mentioned, too, that the disease described in dermatological works as *lepra* (by which we do not mean the true leprosy) is not recognized at all in the present statistics, but is included under the head of psoriasis, and that no attempt has been made to separate the various forms of the latter, described as *P. nummularis*, *P. circinata* (*lepra*), *P. gyrata*, &c., as these are but stages of one and the same disease. There were very numerous illustrations, not only of its tendency to recurrence, but likewise of its markedly hereditary character; and no corroboration was afforded of the view promulgated by Wilson, that psoriasis is remotely transmitted syphilis. One case, how-

ever, was observed in which psoriasis occurred in a lad, with pallid complexion, prominent brow, sunken nose, notched teeth, and other typical symptoms of hereditary syphilis; but it is more than probable that this was a mere coincidence, else, among upwards of 800 cases of psoriasis, such illustrations would probably have been met with more frequently. The local treatment of psoriasis consisted in the use of soothing ointments and lotions in the acute, and of mercurial ointments in the chronic, stage. If there was much irritation of the skin, various kinds of tarry ointments and the liquor carbonis detergens were used, while, in very obstinate cases, which resisted every other kind of treatment, under-dresses of vulcanized india-rubber were sometimes used with excellent effect. The constitutional treatment was, as a rule, most relied upon, especially tonic aperients and alkalies in acute cases; and arsenic, carbonate of ammonia, carbolic acid, and tar (especially coal-tar) in the chronic. It has been well remarked that the subjects of this complaint are usually remarkable for their healthy appear-

ance ; but notwithstanding, an inquiry into the family history often elicits a hereditary tendency to phthisis, and in a few cases the patients themselves gave evidence of the presence of strumous disease : in these, cod-liver oil, steel, and such-like remedies, yielded good results.

Under the head of *Acne vulgaris*, by which is meant inflammation occurring around the orifices of the sebaceous glands as the result of the accumulation in them of hardened and therefore irritating sebaceous matter, which appears on the surface as little black specks, 342 cases were registered, 288 of which occurred amongst the hospital patients, and 54 amongst the private ones. This disease most frequently made its appearance about the period of puberty, and in females often became aggravated at the menstrual periods. It is a most obstinate affection, and little impression was as a rule made upon it by constitutional treatment, even arsenic being generally unavailing. But in those exceptional cases in which the tendency to the formation of little abscesses, the skin covering which was violet, was well marked, cod-liver oil and steel and

strengthening treatment generally was of great use. The local treatment usually consisted in squeezing the little hardened masses of sebaceous matter (comedones, as they are termed) out of the glands, the lancing of pustules and tubercles, and the use of local stimulants, such as lotions of perchloride of mercury and sulphur. A very elegant formula for the latter was recommended to me by Dr. Bulkley, of New York, and which is as follows:— Sulphate of zinc and sulphuret of potassium, of each 1 drachm ; rose-water, 4 ounces : shake the bottle ; cover the affected parts with pieces of lint dipped in the lotion, which are to be removed when the lint is dry. To be used night and morning.

The disease marked Rosacea in the classified list is usually spoken of as acne rosacea, owing to its being frequently associated with an eruption resembling the spots of acne ; but the typical disease is a distinct affection. The first stage in the morbid process consists in dilatation of the capillary blood-vessels, which is followed by congestion and inflammation of the skin, and, if the disease is not arrested, in the new formation of cellular

tissue, so that ultimately tubercles, and even tumours, are developed. It was much more frequently met with amongst the private than amongst the hospital cases, probably on account of the greater annoyance which the disfigurement gave rise to in the former, 21 cases having occurred amongst the 1,000 private, and only 37 amongst the 10,000 hospital patients. It was often the result of intemperance, especially in the male sex, but much more frequently of debility; hence the constitutional treatment, which proved so comparatively useless in *acne vulgaris*, was the *sine quâ non* in *rosacea*. Stimulating local applications were likewise of the greatest service, such as the rumex ointment mixed with sulphur in the proportion of one to two drachms to the ounce. The following is the formula for the preparation of the rumex ointment:—Rumex root, nine ounces; lard, six ounces; yellow wax, one ounce; water, sufficient quantity. Wash and bruise the roots; boil for two hours and strain; evaporate to four ounces; add gradually to the wax and lard previously melted, and keep stirring until cold.

Of the three diseases which alone are capable of implicating the whole body, pemphigus, pityriasis rubra, and lichen ruber, the first was most frequently observed, 16 cases having been met with, one of which was an instance of that rare form described as pemphigus foliaceus. The other two complaints occurred seven times and once respectively. From this it appears that the true lichen ruber, as pictured by Hebra, is a much rarer complaint in this country than in Austria, although the milder forms, described by Wilson under the name of lichen planus, and hitherto classed under the head of Lichen, are by no means uncommon. The case of lichen ruber above referred to occurred in a farmer, about sixty years of age, who consulted me twice in August 1868. The disease had then existed for three years, and implicated the entire cutaneous envelope. At that time he was treated, with slight improvement, by means of tar capsules, and a lotion of liquor carbonis detergens. I saw him again twice in consultation with Dr. Skene of Helensburgh, in April and May 1870. The skin then appeared to be in the same state as when I

saw him first, but he was thinner, and complained much of debility. Small doses of iron and arsenic were prescribed, and an under-dress of vulcanized india-rubber was recommended to be worn. I have not seen him since, but I understand that he is still alive, though gradually losing ground.

CHAPTER IV.

THE second class of diseases not defined by uniform causes, the New Formations, includes as many forms of disease as the Inflammations; but the number of cases was much less, being 251, of which 218 occurred amongst the 10,000 hospital and 33 amongst the 1,000 private patients.

Under the head of Varix 45 cases were registered. But this does not at all represent the actual number, for those complicated with ulceration, erythema, eczema, and syphilis were classed under these heads.

Of Verruca, or Warts, there were 33 examples, which included several cases in which numerous warts had developed on different parts of the head only, or less frequently in combination with warts on the hands and other parts. Isolated

warts were treated locally by means of caustics, excision, &c.; but when numerous, constitutional treatment was more relied upon, such as attention to the general health, tonics, and, above all, arsenic, which in some cases caused their complete disappearance. This leads me to remark that it is a mistake to suppose that warts are mere local affections; indeed, I have little doubt that they are remotely connected with the scrofulous diathesis—that they are what we may call, in imitation of our friends across the Channel, “a benign scrofulide,” as they are specially apt to occur in persons of a scrofulous habit, and as strumous affections very often take on the warty character. This is well illustrated by that form of strumous disease which I have named *Scrofuloderma verrucosa*, and to which allusion has already been made.

Only two cases of *Scleroderma* came under observation, one of which was a hospital case and the other a private patient. The following are the main features of the latter. Janet A—, shopkeeper, came for advice on Jan. 1st, 1867.

Her general health was not quite up to the mark; her tongue was very white and chapped, and the papillæ prominent. Her appetite was extremely capricious, her bowels very costive, and she slept badly. The disease set in about the beginning of February, and was attributed to her having caught cold during the frosty weather. It implicated both arms, the right much more than the left, but did not extend beyond the shoulders. The skin, in irregular patches, was deeply pigmented, especially on the back of the hand and along the outside of the forearm. The whole arm looked a little thicker than natural, and the skin was tense and glossy. Its natural softness, too, had given place to a firm doughy feel, but without any trace of œdema. The forearm was much more affected than the upper arm, but the hypertrophy was by no means limited to the brown parts. Owing to the rigidity, the hand could only be partially closed. The patient complained of neither pain nor uneasiness of any kind. Notwithstanding the use of tonic-aperients and of tonics, the disease had

made some progress when she was again seen on Aug. 22nd, and there was a tendency at some parts to eczematous complication. On Oct. 12th, however, having just returned from a visit of a month's duration at Arran, she came to me looking much healthier, and feeling "a great deal better in herself." The tendency to eczematous complication had disappeared. The arms, especially the upper arms, were softer, the pigmentation paler, and she could close her right hand a little better than previously. Since that time I have not heard anything of her.

Eight cases of Elephantiasis Arabum came under observation, only one of which occurred amongst the private patients. The most typical of these was a poor girl who came for advice on June 22nd, 1866, and whose symptoms were in great measure removed by the tying of the external iliac artery by my colleague, Dr. George Buchanan. As, however, this case was very fully described and illustrated in the *Journal of Cutaneous Medicine*, vol. i. p. 180, it is unnecessary to dwell upon it; but it is only right to add

that about a year after the operation the disease was reported to be again in the ascendant.

Only two cases of Elephantiasis Græcorum (true leprosy) were noted, one of which occurred amongst the hospital, and one amongst the private patients. Indeed, only one other case has occurred in my own practice, showing the extreme rarity of the affection in Scotland. The hospital patient was a young Highlander who had never been out of the country. One of the private patients was a boy aged twelve, who returned from India four years ago, a year after which the symptoms in a mild form began to show themselves. He exhibited brownish-red, circumscribed, erythematous patches of varying shape and size, and a few tubercles, especially on the face, arms, and legs, and accompanied by considerable swelling of the hands. He was thin, and did not look very strong. The other private patient was a young lady aged fifteen, sent to me by Dr. Moir, of Edinburgh. She came home from Jamaica at the age of seven, some time after which the disease commenced on the face,

The eruption consisted for the most part of yellowish-red, erythematous patches, healing in the centre and spreading at the edges, so that elevated rings and segments of circles were left, some of them of very small size, others having an area nearly equal to that of the palm. Here and there the elevation of the eruption was decided, and tubercular in character. The sensation was much deadened, especially where the eruption was most elevated. The parts which had previously been affected were not cicatricial, but sensation had not returned to them. She complained of partial loss of power in the upper extremities, and a large round excavated ulcer was observed in the sole of one foot, which had been preceded by the development of a bulla. This ulcer was situated in the middle of a large erythematous patch upon the sole. Her general health appeared to be good. A course of tonic treatment was, however, recommended. It was also advised that the patches should be dressed with pieces of lint spread with mercurial plaster. I have had no report of her since.

Only one illustration of primary cancer of the skin came under my observation. This was the case of a lady aged about sixty, who was sent to me on July 15th, 1867, by Dr. Francis Henderson, of Helensburgh. She had never been strong, but had had no serious illness. She was of a very spare habit of body, and looked rather younger than she was. She complained occasionally of feeling bilious, and her bowels were rather loose and very easily and powerfully affected by aperients. The skin affection, which was of eight months' duration, commenced over the left breast, and gradually extended over the greater portion of the front of the chest and upper part of the abdomen. There were very slight indications of disease on the back and arms, while the face, neck, lower extremities, and lower portion of the trunk were unaffected. The skin affection presented three elements—(1) a rosy erythema, which appeared to be the first stage; (2) purple streaks, the colour of which was not altered by pressure, which were evidently dilated and obliterated cutaneous vessels, and which seemed to

be the second stage ; and (3) little firm nodules varying in size from pin-heads to split peas, some of them elevated, most of them not, and which were adherent to the cuticle, which was of the natural colour, except over a few of them, where it was rosy. The patient complained of slight itching, and for about three weeks previous to my seeing her she had experienced once or twice a day "a pain shooting up from the right breast." On the 2nd of March, 1868, Dr. Henderson wrote to me as follows:—"The lady I sent to you in July or August with tuberculous cancer of the skin is still alive. The disease has spread over most of the skin of the chest and abdomen, and there are hard tuberculous nodules on the skin of the arms. She suffers no pain, but is now fast sinking."

Only one case of multiple fatty tumours is noted in the present statistics—that of a gentleman, aged about fifty, who resided in the Highlands, whose general health appeared to be good, except that he was the subject of psoriasis, and frequently noticed that his urine was muddy.

He had accidentally noticed the tumours a short time previous to my seeing him. They were scattered over the body, especially the extremities; they were seated in the subcutaneous cellular tissue, varying in size from a large bean to a small walnut; and were soft, nodulated, and painless. They were non-adherent; and the skin over them was of the natural colour. No cause could be assigned for their appearance. A very similar case, not included in the present statistics, was recently admitted into the medical wards of the infirmary, under my care; but in this case the tumours, which presented all the characters above indicated, were associated with the diffuse development of fatty matter in the subcutaneous cellular tissue in the right hypochondriac and epigastric regions, to such an extent that he was obliged to bring the opposite sides of his waistcoat and trousers together with the aid of pieces of string.*

The last class of diseases—the hæmorrhages—

* This case is fully reported in the *Glasgow Medical Journal* for August 1872.

includes a smaller number of cases than any of the others, not because they are infrequent, but because they are not usually regarded as diseases of the skin. Only nine cases were observed, all of which were varieties of purpura, although the case of ephidrosis cruenta, alluded to under the head of "functional affections," ought, strictly speaking, to have been classed with the hæmorrhages. It will be noticed that two cases, one of them a hospital and the other a private case, of that variety of purpura described under the name of purpura rheumatica (peliosis rheumatica—Schönlein) or purpura pemphigoïdes, occurred; and it is worthy of remark that, while in both these cases the bullous and the rheumatic elements were present, in some, one or the other is absent. The following is the note taken of the hospital case:—

Martha B. M——, aged twenty-four, came for advice at the Dispensary for Skin Diseases on Jan. 13th, 1869, on account of an affection of the lower extremities of three months' duration. At that time round red spots, for the most part the size of a fourpenny-piece, made their appearance :

the next day they assumed a bluish tint ; and on the third disappeared, leaving yellowish stains. On the fourth day a fresh crop was observed, which ran the same course, and was succeeded by successive outbreaks up to the time of her coming to the dispensary. When these symptoms first appeared they were accompanied by great swelling of the ankles, with severe aching pain in them. At the same time a large bulla, about the size of a halfpenny, formed on the outer side of the right ankle. About a fortnight previous to admission a similar bulla was detected on the inside of the same leg. On examination, the remains of this bulla were still apparent ; and numerous purpuric spots, varying in size from a pea to a sixpence, and of a reddish, livid, or yellowish tint, were scattered over both lower extremities. The veins were not varicose. She seemed in pretty good health ; her gums were healthy ; she had always had plenty of food of good quality, and had never before suffered from rheumatism. The treatment consisted of ten drops of turpentine on sugar three times a day, with occasional doses of

castor-oil, which had an immediate effect upon the disease : but it was two months before all tendency to the development of purpuric spots had disappeared, and for some time she was troubled with an ulcer which formed in the site of the last bulla as the result of injudicious local applications.

The following is an abstract of the private case, which was seen by me in consultation with Dr. S. J. Moore, of Glasgow, on December 9th, 1868. The patient was a stout, healthy-looking gentleman, who "had never had a day's illness in his life," but who was interested in, and probably excited by, the municipal elections which at that time were pending. For several weeks he complained of pains, or rather perhaps of weakness, in his joints, especially those of the lower extremities ; but his illness, which was accompanied by febrile symptoms, set in decidedly only a few days before I saw him. His tongue was moist and not much coated ; his appetite fair ; his bowels had been freely opened by medicine ; and his urine, which previously had been scanty and loaded with lithates, was clear and of fair amount, under the

influence of acetate of potash and colchicum. His pulse was 120 and rather feeble, but its rapidity was out of proportion to the extent of the fever. The lower extremities were markedly œdematous. There was swelling around some of the joints, and a feeling of stiffness rather than of pain in them, which was attributed to the swelling. Three morbid elements were observed on the skin—namely, purpurous spots, a papular eruption, and bullæ. The first varied in size from mere points to ecchymoses the size of the palm and for the most part round. The greater number of the small ones constituted apparently the second stage of the papular eruption; and on the back, where the latter was most abundant, the transition from papules to purpurous spots was observed—that is to say, papules whose elevation was subsiding, and whose redness only partially disappeared on pressure. Most if not all of the large ecchymoses occupied the seats of previous bullæ, which varied in size from a split pea to a goose's egg, and which, when at their height, were fully distended with clear serum. The black purpuric patches covered

by the flaccid envelopes of the bullæ after the serum had escaped looked very like patches of skin which had mortified. The bullæ were most abundant on the legs and buttocks; and the cutaneous manifestations had no special tendency, such as is often observed, to implicate the skin around the joints. The acetate of potash was continued, and purgative doses of castor-oil and turpentine recommended, along with light, nourishing, fluid food. This patient made a good recovery.

Although not included in the present statistics, the following case, which was seen by me in consultation with Dr. Adam of Coatbridge while these papers were being written, is worthy of note, owing to the rapidity with which the disease proved fatal:—A mining engineer, aged twenty-seven, who had previously enjoyed good health but who had lost a brother of phthisis and a sister of “softening of the brain,” began to complain of debility, “weak stomach,” and constipation, as the result, it was supposed, of hard work, and irregularity as to his meals. His symptoms did not improve under treatment; and he

was therefore sent to Arran for change of air. On Friday the 16th of August, while there, his eye became ecchymosed, when he immediately returned home. When I saw him on Monday the 19th, copious extravasation of blood was observed beneath the conjunctivæ and around the eyes, while small purpuric spots were detected upon the legs, which had only appeared on the morning of my visit. His gums were bleeding, he had slight hæmorrhage from the stomach and bowels, and the urine was bloody. Two days afterwards he died. Dr. Adam wrote me as follows with regard to him:—"After he got the castor-oil and turpentine the hæmorrhage from the kidneys and mouth became much less; but it never altogether ceased from the stomach till a few hours before death, up to which time the retching and vomiting of altered blood were very persistent. Although stimulants were given very freely, exhaustion came gradually on, accompanied with delirium." I regret that in this case the subcutaneous injection of ergotine, which was spoken of, was not tried.

The last case,—which is named *Purpura tuberculosa* in the classified list, but which might with equal propriety be termed *P. verrucosa*, or, as suggested by my late colleague, Dr. A. B. Buchanan, who saw it along with me, *P. hypertrophica*,—was that of a woman aged sixty-six, who had formerly been in service, whose family history was good, and who had previously enjoyed excellent health. Four months prior to my visit, as the result, it was supposed, of poor fare, she began to complain of pain and swelling of the right foot; shortly after which slightly elevated livid spots, about the size of pin-heads, made their appearance on the thigh. These rapidly increased in number and in size until they became as large and as elevated as split-peas, especially on the inner and posterior aspects of the thigh. In these situations the surface of the patches, which were blackish, gradually assumed a distinctly warty appearance and feel. At this time, too, the whole limb became thicker, so that its diameter was about a half greater than that of the left. This increase was, in the foot, prin-

cipally the result of œdema ; but in the leg and thigh it was mainly, if not entirely, due to induration of the cellular tissue. When I saw the patient this induration was not limited to the limb, but implicated likewise the right side of the abdominal parietes and the right breast, which was firm, round, and plump, while the left was flabby and about one-third the size. The left thigh was at this time affected similarly to the right, but to a much slighter extent. The legs were quite free of eruption. The patient complained of neither pain, itching, nor heat ; but on placing the hand upon the eruption, especially upon the warty part, it felt much warmer than the healthy skin. (The temperature was one degree higher.) The right limb felt numb and weak, and she was confined to bed owing to the uneasiness experienced in sitting. She was thin, but not more so than many people of her age ; her appetite was bad, and her bowels exceedingly costive, as the result, probably, of the confinement in bed. Otherwise she appeared well, and was very cheery. She

gradually sank, three months after I first saw her, and seven months from the commencement of the disease, suppuration having occurred shortly before death at those parts which were in contact with the bed.

PART II.

*THE THERAPEUTICS OF DISEASES OF
THE SKIN.*

THE THERAPEUTICS OF DISEASES OF THE SKIN.

CHAPTER I.

IN referring to the therapeutics of diseases of the skin, I shall not content myself with a mere enumeration of the remedies which are likely to prove of service, as is so often done—for such a course is practically of little use; nor, on the other hand, shall I attempt an exhaustive account of them: my remarks are intended to be suggestive only, and I must leave the filling in of the picture to the subsequent experience of the reader.

A great deal has been written with regard to the danger of suddenly “driving in” an eruption; but while I am not prepared to deny that its

sudden disappearance is in no instance prejudicial, I am quite convinced that there is very little reason to dread such an occurrence. The idea has in part arisen from the fact that an eruption of the skin generally disappears if the patient is seized with any severe internal malady, in which case the result is mistaken for the cause. Some time ago—and such cases are frequently met with—while in attendance upon two children for very severe attacks of chronic eczema affecting the greater portion of their bodies, one of them took measles, and two or three days thereafter the eczematous eruption had almost disappeared. In a few days the other child fell ill of the same complaint, and in her case, too, the eczematous eruption quickly vanished. There could be no doubt that these children were affected with measles, for, while it was difficult to make out the characteristic eruption on their skins, owing to the existing eczema, their younger brother presented about the same time all the characteristic symptoms of measles. When the disease had run its course

in the case of the two first-named children, the eczematous eruption gradually but perseveringly returned—a circumstance which unfortunately happens in the great majority of cases. Had the intercurrent disease been an inflammation of an internal organ, instead of a specific fever, its occurrence would probably have been attributed to the disappearance of the eczema. Those who are most intimately acquainted with the treatment of diseases of the skin, will probably agree with me in thinking that the difficulty lies, not in curing them sufficiently slowly, but in curing them with sufficient rapidity.

In the treatment of diseases of the skin, as of most other complaints, it will be found that there are often different ways of arriving at the same goal. In Germany, especially in Vienna, local treatment is principally relied upon, while in England it appears to me that constitutional treatment is too exclusively resorted to. Some diseases of the skin are best treated by local, others by constitutional treatment, while a few may be cured by either; but the majority of

them yield most readily, and are removed most permanently, by a mixed local and constitutional treatment.

A.—LOCAL TREATMENT.

Local treatment is *especially applicable to eruptions dependent upon local causes*—is, indeed, the only treatment, in most cases, which is of any use. In scabies, for example—*i.e.* the disease due to the presence of the itch insect—external treatment suffices for a cure, it being unnecessary, as suggested by Wilson¹ and others, to prescribe sulphur internally. Frequently, however, as in cases of eczema, an eruption is called forth by a local cause, which would have failed in producing such a result had there not been some constitutional derangement. Such cases require a mixed treatment.

On the other hand, it must be borne in mind that *many eruptions originally dependent upon constitutional causes become ultimately mere diseases*

¹ On Diseases of the Skin. Sixth Edition, pp. 223. London; Churchill, 1867.

of the skin, the constitutional cause having disappeared, and the eruption being perpetuated owing to the skin having, so to speak, contracted a bad habit. We see this, for instance, occasionally in young adults affected with chronic eczema which had begun in infancy; and in such cases a local treatment often yields the most satisfactory results.

We must not, however, lose sight of the fact that *local applications have not a merely local action*—that they are absorbed, and often react on the system at large. In the case of mercury this is admitted and acted upon; in the case of many other drugs it may be admitted in the abstract, but it is seldom taken into account. For instance, it occasionally happens that when a preparation of tar is rubbed firmly over an extensive surface, indications of its absorption are not wanting. The patient may be seized with nausea, vomiting, and diarrhoea, the discharges from the stomach and bowels having a black colour, while the urine may have a dark-green or even blackish tint, and a tarry odour.

It is not usual for all these symptoms to be present in a given case, but the occurrence of any of them is sufficient to show that the remedy is not a mere local one.

Lastly, we must bear in mind that *local applications are far from uniform in their action*; this being due in part, no doubt, to the state of the general health, the varying sensibility of the skin, and the degree of care with which they are prepared. It is of special importance to ensure the careful preparation of ointments; for if they are the least gritty or rancid they are very apt to prove injurious.

If the affected surface is covered with scales or crusts, or *débris* of any kind, it is generally desirable to remove them at the outset. This is often necessary for purposes of diagnosis, for otherwise we can only guess at the condition of the parts beneath: it is still more important previous to treatment, for otherwise local applications cannot reach the diseased surface whose condition they are intended to modify. The removal of crusts is a very easy matter, and may

be effected in various ways. I usually order the patient to saturate the parts thoroughly with almond-oil, and to remove the crusts thus softened with warm water. Should this fail, a poultice of crumb of bread and hot almond-oil may be applied at night, and if they do not come away with the poultice in the morning the part may be lubricated with fresh almond-oil, and the crusts removed about half an hour afterwards with the finger-nail, or, if the disease is on hairy parts, with the comb. The removal of scales — in cases of psoriasis, for example—is often satisfactorily accomplished by the use of potash applications, to which reference will be made hereafter.

Leaving out of view for the present the exact nature of the eruption (*i.e.* whether syphilitic, simple, strumous, parasitic, or the like), the first point to decide, before treating an inflammation of the skin, is *whether the eruption is acute or chronic*; that is to say, whether it requires soothing, or will tolerate stimulating applications.

If the surface is acutely inflamed, if it is the

seat of a copious eruption of vesicles or pustules, if there is much swelling, and, above all, if the patient complains of burning heat or pain rather than of itching, local treatment should either be avoided altogether or soothing applications resorted to. To this as to most rules, however, there are a few exceptions. In tinea sycosis (ringworm of the beard), for instance, when the part is greatly swollen and indurated, the seat of numerous pustules, and the source of burning heat and pain, the best treatment consists, not in the use of soothing applications, but in the extraction of the hairs. In this case the treatment is decidedly stimulating; but the stimulation is more than counterbalanced by the soothing effect of removing hairs which are diseased, which are loaded with the spores and tubes of the fungus, and which are acting as foreign bodies.

Soothing applications may take the form either of baths, powders, poultices, ointments, or lotions.

Baths are often serviceable, especially warm baths to which a little washing soda, a pound of gelatine, or a few handfuls of starch have

been added. They are particularly useful in inflammations affecting a large extent of surface; in cases of acute general psoriasis, for example.

Benefit is sometimes derived, too, from *hydro-pathic treatment*, which I am in the habit of administering thus: "On a firm mattress a sheet of M'Ghie's oil-paper or other material, to prevent the wetting of the mattress, is placed; then a couple of straps are laid across the bed; above this a dry blanket is spread out; and, finally, a sheet wrung out of cold water. The patient in a state of nudity is made to lie down on this; and it is then wrapped tightly and carefully round him, so that every part of the skin is, if possible, in contact with it, and the blanket is similarly applied. The whole is then kept in position by the two straps, and the patient is covered by two or three blankets thrown loosely over him. In this state he is allowed to remain for two or three hours, during which time he is allowed to drink water *ad libitum* to promote perspiration; and the process is completed by making him

jump into a cold bath, and then dress himself and take a sharp walk." This treatment is of value in acute general psoriasis associated with great heat of skin; it is sometimes useful—nay, even curative—in chronic general psoriasis, and relieves irritation of the skin when present in a very marked manner. It may, therefore, be repeated daily, and continued as long as improvement takes place.

The *shower bath* sometimes yields good results. It may even remove the eruption altogether, but it is chiefly of benefit as a palliative for the relief of uneasiness—in severe cases of acute and chronic eczema, for example. The water should be finely divided—should fall from the height of about a foot, and the shower should be continued on each occasion from five to fifteen minutes. The bath may be repeated three or four times a day, and the patient should take brisk exercise for half an hour after each.¹

Before leaving the subject of the treatment

¹ On Diseases of the Skin, including the Exanthemata. By Ferdinand Hebra, M.D. Vol. ii. p. 147. Sydenham Society's Translation.

of diseases of the skin by water it may be well to state that patients should be warned against the use of hard or of sea-water, which is very apt to bring out, or to aggravate, an existing inflammation. Spring water is generally hard, and therefore to be avoided, while distilled, or rain water, is always to be preferred when it can be had; if not, the water should be boiled so as to cause a deposit of its salts before it is used.

Absorbent powders are sometimes of use when we have to deal with an acute inflammation, such as erysipelas or shingles, and also when there is a tendency to moisture, and when it is desirable to keep the parts dry, as when we have to do with an eczematous eruption implicating portions of skin which are in apposition. Those which I am most in the habit of using are: Powdered starch, zinc, lycopodium, talc, carbonate of magnesia, and carbonate of zinc. These may be combined in various ways, and when burning heat or itching is complained of, a little camphor may be added, as in the follow-

ing prescription: Powdered starch, six drachms; oxide of zinc, three drachms; cochineal, one grain; powdered camphor, half a drachm: make into a powder. Dusting powder, to be kept in a stoppered bottle.

Poultices are often of value for the relief of pain or tension, or to hasten suppuration when it is impending; but it must never be forgotten that their long-continued use is a source of inflammation. Thus they are apt to call forth eczematous eruptions, especially in persons pre-disposed thereto; and most persons are familiar with the fact that their incautious application in the treatment of boils favours the development of others in the vicinity. The most soothing poultices with which I am acquainted are made with cold water and potato starch.

Soothing ointments are more universally applicable than any of the preceding, although, in some cases, even the most emollient and most carefully prepared, owing to some idiosyncrasy, cannot be tolerated. They are used to soften and favour the removal of crusts and other *débris*,

but, above all, to form a covering for and protection to the inflamed parts, and thus to exclude the air. One of the best of these is the benzoated oxide of zinc ointment (so prominently brought before the profession and so justly lauded by Wilson), and which is much improved by the addition of two drachms of spirit of camphor to each ounce of the ointment, being thus rendered softer and at the same time more cooling. The benzoin in the zinc ointment prevents it from becoming rancid and irritating, while at the same time it imparts to it a certain fragrance.

Another very soothing ointment is one containing bismuth, as in the following mixture: Sub-nitrate of bismuth, half a drachm; rectified spirit, a drachm and a half; simple ointment, six drachms; oil of roses, half a minim. We must avoid using benzoated lard in this prescription, for while the benzoin in the zinc ointment, for reasons which I cannot explain, does not irritate, it is apt to do so in some persons when combined with other medicaments, such as bismuth.

A very good ointment, too, though not so soothing, is made from litharge plaster and olive-oil, as follows: Litharge plaster, four drachms; best olive-oil, three drachms. Apply heat until the plaster is melted, then stir till the mixture cools. This ointment, spread on strips of linen and applied in the form of a "scultetus," is very soothing, and gives support to the parts in cases of acute eczema of the legs. In cases of strumous eczema such as often occurs on the hairy portions of the face, and of strumous ulceration, great benefit is frequently obtained by keeping the parts saturated with pure cod-liver oil.

Soothing lotions are sometimes of service for the relief of uneasiness; indeed they are not unfrequently curative—in cases of acute eczema impetiginodes of the face, for example. Those containing lead and soda are amongst the best, and may be used in the following forms: (a) Solution of subacetate of lead, one drachm; glycerine, four drachms; distilled water, six ounces. (b) Dilute hydrocyanic acid, two drachms;

bicarbonate of soda, one drachm; glycerine, four drachms; rose-water, five ounces and a half. The parts should be frequently sponged with these lotions, or pieces of clean rag dipped in them may be kept constantly applied.

CHAPTER II.

WE have seen that in the local treatment of diseases of the skin we must take into account whether the eruption is acute or chronic. In the last chapter we considered that which is applicable to acute inflammations, and now we have to study that which is applicable when the eruption is chronic.

If itching is a marked feature (and, for the present, we leave out of consideration diseases dependent on parasites, such as scabies—the itch—which require special treatment), we may expect to derive benefit from the use of *emphyreumatic oils or their derivatives*. It is right to mention, however, that, in exceptional cases, these increase the irritation to an intolerable

extent ; but in the majority they not only relieve the itching, but are decidedly curative, especially in chronic erythema, chronic eczema, and psoriasis (although more slowly), while in prurigo they give immense relief. It is well to know that in some persons, with very sensitive skins, their application produces all the phenomena of dermatitis (burning heat, redness, swelling, and sometimes an eruption of vesicles and bullæ). Their long-continued use, too, produces, in all persons, though more speedily in some than in others, an inflammation at the orifices of the hair-follicles : papules and pustules make their appearance, which present this peculiarity, that in the centre of each spot a black, tarry point is observed. This eruption has been called *tar acne* by Hebra. Some time ago I prescribed for a little girl, who was affected with slight chronic eczema of the leg, a lotion containing one of the empyreumatic oils. About six weeks afterwards her mother brought her to me again, the lotion having been continued regularly in the interval, and informed me that the disease was a great deal worse. On

examining the part, I found that the eczematous eruption had quite disappeared, and that its place was taken by an eruption of tar acne. The lotion was discontinued, and the artificial rash rapidly subsided.

The empyreumatic oils with which I am most familiar are—first, *Pix liquida*, or common Wood-tar, of which Guyot's solution of tar is a purified form, and possesses these advantages, that it mixes with water in all proportions, does not emit such a pungent odour, and does not discolour the skin to the same extent; second, Coal-tar (*Pix mineralis*), of which the best form is the *liquor carbonis detergens* of Wright and Co., a solution which forms a creamy emulsion with water; third, Oil of Cade (*Oleum Cadini*), which is the product of the dry distillation of the wood of the *Juniperus oxycedrus*; fourth, the *Oleum Rusci*, which is obtained from the bark of the *Betula alba*, or white birch. The odour of the last two is not so pungent or disagreeable as that of the others, but they are more expensive, and therefore less adapted for hospital use.

These preparations may be employed alone—in cases of chronic psoriasis, for example, after the removal of the silvery scales—in which case they should be rubbed into the affected parts, twice a day, as firmly as possible short of causing pain, or as lotions or ointments, as in the following prescriptions:—Guyot's solution of tar, soft soap, rectified spirit, of each one ounce; spirit of rosemary, one drachm; mix. Wash off before each re-application. — Nitric oxide of mercury, one scruple; nitrate of mercury ointment, one drachm; oil of white birch bark,¹ one drachm and a half; benzoated lard, four drachms: mix.

Two other remedies of this class must be mentioned—namely, Creasote and Carbolic acid.

Creasote may be added to ointments in the proportion of from five to ten grains to the ounce, for the purpose of allaying irritation of the skin. In cases of chronic psoriasis a strong creasote ointment was recently recommended by Mr. Balmanno Squire, and I have found it useful,

¹ Mr. Greig, the manager of the New Apothecaries' Company, Glassford Street, Glasgow, has obtained a supply of this from Germany for me.

although sometimes too irritating. The following is the form of ointment he employs:—Creasote, two ounces; white wax, one ounce: dissolve. Rub very firmly into the eruption night and morning after the scales are removed.

Carbolic acid has a less disagreeable odour than the tarry preparations, and is more cleanly. It is soluble in water with the aid of a little glycerine or spirit, and forms a colourless watery solution. For these reasons it can often be used when tar is inadmissible, as when an eruption is situated upon an exposed or hairy part; but it must be admitted that, as a rule, it is not nearly so effectual as the empyreumatic oils. The following is a form in which I frequently prescribe it:—Crystallized carbolic acid, two drachms; Price's glycerine, six drachms; rectified spirit, four ounces; distilled water, one ounce: dissolve. Sponge the affected parts two or three times a day, and also when itching is complained of. This solution is of use in cases of chronic erythema, chronic eczema, and the like, and not only relieves irritation of the skin, but also is directly curative.

Potash applications have of recent years been brought prominently under the notice of the profession by Hebra. They are of use in the removal of abnormal products—*e.g.* the scales of psoriasis,—and in many cases they favour the cure of the eruption by virtue of their stimulating effect, especially in combination with other remedies, such as tar.

Those which I am most in the habit of using are, potash soap (black soap—*sapo mollis*), liquor potassæ, and solutions of potassa fusa varying in strength from two grains to a drachm to the ounce of the excipient. I shall refer to stronger solutions under the head of Caustics. A few illustrations may be given of their use.

In cases of *tinea versicolor*, that brown scurfy eruption which attacks the covered parts of the body, due to the presence of a fungous growth, the *Microsporan furfur*, we can effect a cure by scrubbing the *whole* of the eruption night and morning with black soap or its solution in spirit. The soap acts as a stimulant, but is chiefly serviceable owing to its mechanically removing the

scales which are loaded with the fungous matter. But the cure may be accelerated by the use of one of the so-called parasitocides, which can be combined with the solution of black soap as follows:—Perchloride of mercury, twelve grains; black soap, two ounces; rectified spirit, four ounces: dissolve, and add oil of citronella, one drachm. Rub, night and morning, as firmly as possible into the whole eruption short of causing pain.

Again, potash applications are useful in cases of chronic eczema, not only by removing morbid products and allaying irritation, but also by stimulating the parts and favouring the disappearance of infiltration of the skin when it is marked. For this purpose I generally employ solutions of potassa fusa, and vary the strength of the solution according to the amount of the infiltration and the extent of the disease. It should be rubbed into the affected part with a large paint-brush, or with a small sponge provided with a handle, until decided smarting and excoriation result, when its action may be stopped by washing with water and

applying a soothing ointment. If the surface is left alone after the application of the solution, it is soon covered by an immense number of gelatinous drops, showing the extent to which it has been stimulated. After the effect of the first application has passed off, it may be repeated, and the treatment continued until the infiltration of the skin is gone, when the cure may be accelerated by the use of some other application, such as the oil of cade, or the like.

Lastly, in cases of chronic general psoriasis, Hebra's modification of Pfeuffer's treatment may prove effectual. This consists in rubbing soft soap into the whole of the eruption twice a day for six or eight days, each patch being on one occasion rubbed so hard as to remove the scales entirely, and to cause slight bleeding. During this time, and for a few days thereafter, the patient lies in bed enveloped in blankets, after which he takes a warm bath and dresses himself. The treatment often requires to be repeated, and is only successful in a few cases; for which reasons, and also owing to the pain which it produces, and

the necessity of [confining the patient to bed, it can seldom be carried out in private practice.

Mercurial applications are often of service in simple inflammations, although not to the same extent as the preceding remedies ; while in syphilitic affections they are almost indispensable. If itching is a prominent symptom, they may with advantage be combined with one of the empyreumatic oils. Those which are most valuable are corrosive sublimate, calomel, white precipitate, red oxide, and red and green iodide of mercury, and ointment of nitrate of mercury; not to speak of the well-known yellow and black washes of the Pharmacopœia. A few illustrations of their use as powders, lotions, and ointments may be given.

In cases of syphilitic condylomata, at the anus we shall say, Ricord's treatment is very effectual. This consists in washing the parts twice a day with solution of chlorate of soda, drying them, dusting them with calomel, and separating them with a piece of clean dry lint.

Again, in cases of acne and in chronic erythema of the head (pityriasis capitis, as it is often called),

the following lotion is of use :—Perchloride of mercury, from twelve grains to one scruple ; Price's glycerine, four drachms ; rectified spirit, five ounces ; spirit of rosemary, one drachm ; cochineal, one grain : mix and strain. Dip a piece of sponge or flannel into the mixture, and rub firmly into the part night and morning.

In a recent very able article by my friend Dr. Moriz Kohn, of Vienna, on *Lupus Erythematodes*,¹ the value of *emplastrum hydrargyri* in that affection is referred to ; and I can in a measure corroborate his experience of it in the treatment of this and of allied affections. A piece of lint the size of the part is spread with the melted plaster and applied, and changed once in twenty-four hours.

Lastly, when we have to deal with patches of dry eczema (*eczema siccum*), or of psoriasis, and above all, when we have to treat dry syphilitic eruptions, the following ointment may be used :—Subchloride of mercury, one drachm ; ammoniated mercury ointment, three drachms ; glycerine

¹ Archiv für Dermatologie und Syphilis, Erstes Heft, p. 18. 1869.

(Price's), one drachm; simple ointment to one ounce: mix. Rub firmly into the eruption night and morning.

It must never be forgotten that all kinds of mercurial applications are to a certain extent absorbed, and hence patients must be warned of the possibility of salivation, especially if a considerable extent of surface is being acted upon.

CHAPTER III.

IF we were to put faith in the general verdict, we should have to believe that *sulphur* is the panacea for all skin complaints—a verdict which I, for one, cannot homologate; for the result of my experience is, that, taken over head, it gives rise to more diseases than it cures. It is very apt to produce eczematous eruptions, or to aggravate them if they are already present, and all the more certainly the more nearly the attack approaches an acute one, or the more decided the tendency to suppuration. If an eczematous eruption is produced by the nails of a patient attacked by lice or labouring under the itch, sulphur kills the parasites and removes the source of the irritation, and thus the medicine often gets the credit

of curing eczema when in reality it acts by removing the special cause. As the reader is probably aware, sulphur is the nearly universal remedy for the itch; but I have no hesitation in saying that it is often not the best, for although there is no drug more effectual in destroying the acari, there is none which is more likely to irritate the skin, which is already over-irritated by the nails of the patient. I shall refer to this, however, further on.

In the treatment of chronic obstinate patches of eczema the use of sulphur may do good in virtue of its stimulating properties, although it is generally found that other remedies are more effectual.

But while all this is true, it must be admitted that, when used with discrimination, in certain cases sulphur is of great value. Thus in the treatment of so-called acne rosacea (that eruption which is so frequently observed upon the face as the result of debility or the abuse of stimulants) it is exceedingly useful in the form of ointment, as in the following prescription:—*Rumex* oint-

ment, one ounce;¹ hypochloride of sulphur, two drachms: mix. Rub very firmly into the eruption night and morning, short of producing pain or inflammation. Sulphur is useful also in cases of acne vulgaris (that pimply eruption which occurs so frequently on the faces and backs of young persons), in the form of a lotion, as follows:—Sulphur, and glycerine (Price's), of each six drachms; rectified spirit, six ounces; oil of roses, one minim: mix. Shake the bottle: rub very firmly into the part night and morning with a piece of flannel. Finally, sulphur is of great value in *genuine* cases of prurigo (I do not refer to cases of pruriginous eruption—*i.e.* eruptions brought out by the nails of patients whose skins are itchy), especially in combination with one of the empyreumatic oils, as follows:—Sulphur, one ounce; liquid tar, six drachms; benzoated lard, four ounces: mix. Rub very firmly into the eruption night and morning.

During a visit last summer from Drs. Weisse

¹ For the mode of preparing rumex ointment, see Part I. chapter iii. p. 52.

and Satterlee, of New York, those gentlemen recommended me to try a solution of *acetate of soda*, of the strength of eight grains to the ounce of water, in the treatment of lupus. Having given it a pretty extensive trial, I am prepared to recommend it in the treatment of strumous diseases of the skin. I usually employ a solution of twenty grains in an ounce of water, which, as a rule, does not smart. The affected parts are sponged with it three or four times a day, and rags, dipped in it, are applied at night, and covered with oil-skin. It is very valuable in lupus exedens, and in other strumous affections, in the stage of ulceration. Under its influence the crusts fall off and the ulcers heal, after which the improvement is slower, and then a more speedy cure generally results from substituting some other treatment. It is sometimes of advantage to touch the ulcers with the pure crystals of the acetate of soda; while the injection of the solution into strumous sinuses with the aid of a small glass syringe favours their closure to a marked degree. Such is my experience of the value of this

remedy; but if I am asked how it acts, I must confess my ignorance.

We are indebted to Hardy, of the St. Louis Hospital, for the proposal to employ *impermeable dressings* in the treatment of certain diseases of the skin. This treatment has been tried on an extensive scale by Hebra of Vienna and by myself, and the following is the result of my experience of it. The material which is used for the purpose has been, for the most part, vulcanized india-rubber and vulcanized india-rubber cloth. It acts in a variety of ways: it excludes the air, keeps the part warm and at a uniform temperature, and promotes the secretions from the cutaneous glands, which it retains, so that they macerate and favour the removal of the epidermis. According to Hebra, it does good also in virtue of the sulphur which it contains. Of this, however, I am doubtful. I lately recommended a patient labouring under eczema rimosum (the fissured variety of eczema) of the palms to use a pair of vulcanized india-rubber gloves. He subjected the left hand only, however, to the treat-

ment. "For the right," he wrote me, "I merely wore an ordinary leather glove with the fingers cut off, and lined in the palm with a bit of oil-silk; this I have found quite sufficient for it, as the slight indications of its being affected have quite disappeared" (*i.e.* in a couple of weeks).

The following cases illustrate the value of impermeable dressings in appropriate cases.

CASE 1. *Psoriasis palmaris*.—Mrs. M——, aged forty-five, monthly nurse, was admitted at the Glasgow Skin Dispensary August 12th, 1868. The disease had appeared about nine years previously, since which time she had hardly ever been free from it. It commenced as a small crack in the palm of the right hand, and gradually the whole of the palm became affected.

Appearances on admission.—The whole of the palmar surface of the right hand was covered with minute silvery points about the size of pin-heads, while numerous fissures traversed the affected surface, which had always been dry, and not in the least degree itchy. At the period of her visit, according to the patient's statement, the

eruption was on the increase. She was ordered a lotion containing tar and spirit, and a tonic aperient mixture.

On Dec. 7th the eruption was only slightly improved; this treatment was therefore discontinued, and she was ordered to procure an india-rubber glove and to wear it constantly for a fortnight.

On the 23rd the eruption had almost entirely disappeared.

Jan. 6th, 1869.—It was observed that there was a little tendency to excoriation on the sides of the fingers, owing to the maceration of the tender epidermis, as the result of the continued use of the glove.

11th.—As the soreness continued, and as the psoriasis had completely disappeared, the use of the glove was omitted, and she was recommended to apply a little benzoated zinc ointment to the tender parts.

On Feb. 8th the artificial irritation had subsided, and, as the primary eruption remained in abeyance, notwithstanding the cessation of the use of the

glove, she was dismissed cured. (Reported by Mr. J. D. Walker.)

CASE 2. *Eczema manuum*. — Allan M'A——, aged forty, hatter, was admitted to the Glasgow Skin Dispensary on Dec. 23rd, 1868. The disease had appeared for the first time about three years before this date, and somewhat in the following manner. He first noticed a number of small "blisters," about the size of pin-heads; these, after remaining a short time, burst, discharging their contents, and leaving a raw surface, which was extremely itchy and "leeted" very much. The eruption appeared first on the dorsal surface of the right hand, then on the sides of the fingers, and lastly on the dorsal surface of the fingers. Shortly after this a similar eruption appeared on the left hand. The disease lasted a considerable time, and disappeared under medical treatment. The second attack commenced about eight weeks previous to admission, appearing in the same manner and order as before; but this time the palmar surfaces of the fingers were implicated as well as the above-mentioned parts.

Appearances on admission.—On the left hand the eruption implicated its dorsal surface, as well as the dorsal surfaces and sides of the fingers: these parts were considerably infiltrated, covered with a serous exudation, and were the seat of numerous excoriations. The right hand presented similar appearances, but the infiltration of the skin on its dorsal surface was much more marked. The eruption was extremely itchy, and was rapidly extending. The patient's general health was good. He was ordered to procure a pair of india-rubber gloves, and to wear them constantly for a fortnight.

On Jan. 6th, 1869, the disease had in most parts completely disappeared, a faint reddish blush only being left on the sites of the previous eruption. He was ordered to continue the use of the gloves for another fortnight, and then to return—which, however, he failed to do. (Reported by Mr. J. D. Walker.)

CASE 3. *Eczema capitis*.—Louisa W—, aged four years and eight months, admitted to the Glasgow Skin Dispensary June 28th, 1869. Her father stated that the eruption first made its

appearance when she was three months old, and disappeared about three months thereafter. She remained well till after an attack of scarlatina at the age of a year and a half, when it reappeared, since which time she had never been altogether free of it.

Appearances on admission.—The eruption implicated the external ears and the whole of the scalp. These parts were very red, much infiltrated, exuded an abundance of clear serum, and were studded with crusts. The patient complained of great irritation of the skin and of burning heat. She looked rather delicate, but her digestive organs were in good order, although the year before she had been troubled with worms. Various remedies had been tried, but to little purpose. She was ordered a vulcanized india-rubber cap, which was to be worn constantly.

On July 15th the infiltration of the skin and the exudation had completely disappeared, the itching and burning heat were almost gone, and a faint red blush was all that remained of the previous eruption. The india-rubber cap was con-

tinued. The patient did not return. (Reported by Mr. Robert Sinclair.)

CASE 4. *Strumous disease of the toes, &c.*—A young lady, aged about twenty, consulted me in 1868 with regard to an attack of lupus, of old standing, implicating the tip of the nose. It was of the non-ulcerating variety. She had also a strumous eruption on each foot on and around the toes, associated with several small ulcerations, and covered for the most part with thick hardened epidermis. Under treatment the nose recovered perfectly, and the eruption on the toes improved, but the skin remained very thick and hard, and would not yield to the remedies employed. She was therefore ordered to cover these parts with vulcanized india-rubber. This was on April 13th, 1869. She made use of a piece of a tobacco-pouch for each foot, and at her next visit, on April 28th, the thick, hardened epidermis had almost entirely disappeared, and the toes had nearly resumed their normal appearance. She was ordered to continue the use of the india-rubber for a considerable time.

CASE 5. *Pruritus senilis*.—A gentleman, aged about sixty-five, of a spare habit of body, whose general health was fair, with the exception of a marked tendency to constipation, and who had the advantage of plenty of out-door exercise, consulted me on account of intolerable itching over the whole body, that malady which is described under the title of *pruritus senilis*, and which has no connection, as some would have us believe, with *phtheiriasis corporis*, the disease due to the presence of lice, and which is so common in old persons. His skin was the seat of a pruriginous eruption, the result of the scratching in which he indulged, and the irritation of the skin interfered much with his sleep. After a trial of other means, which were only very partially successful, he procured at my request a complete underdress of vulcanized india-rubber cloth, on commencing to wear which the itching and eruption completely disappeared.

This treatment is useful also in cases of *prurigo* and *ichthyosis*, and probably in most obstinate localised eruptions. It possesses this advantage,

too, that it does not interfere with the simultaneous use of other local applications. The dressing often proves curative, even when the patient is only able to make use of it at night; but of course it is more quickly and more certainly successful if worn continuously. It should be removed from time to time for the purpose of cleansing it, and of wiping the skin; and if, as often happens, the tender skin in the vicinity of the eruption is abraded and raw, a little benzoated zinc ointment may be applied with advantage, or the covering taken off for a day or two.

CHAPTER IV.

IT is unnecessary to enter into details with regard to the use of *caustics*, as their selection and application must be regulated, for the most part, by the same rules as have been so well laid down in surgical works ; but a few illustrations of their special use in certain diseases of the skin may be given. The use of *arsenical caustics* is not without risk, especially if a considerable extent of surface has to be acted upon, for the danger of their absorption to such an extent as to give rise to symptoms of arsenical poisoning is by no means remote ; indeed, many cases have been recorded—most of them, I am glad to say, in the hands of illiterate persons—in which a fatal issue has resulted in this way. It is right to add, however, that in some cases of epithelioma, associated with ulceration of small

extent, I have sometimes succeeded in healing up the ulcers by painting them daily with *Fowler's solution*—taking care, however, that not more than from ten to twenty drops of the solution are used on each occasion. This treatment is by no means so free from pain as might be expected.

In cases of chronic eczema associated with very marked infiltration of the skin, and when the eruption is not very extensive, the application of a solution of *potassa fusa*, in the proportion of one drachm to two of water, as recommended by Hebra, is of great value. The affected part is painted with this solution with the aid of a large brush, after which the patient is directed to dip the hand or a piece of flannel in water and to rub the skin until a lather is produced. Cold-water dressings are then applied, changed frequently, and continued till the effect of the first application has passed off. It may be repeated as often as necessary until the infiltration of the skin has been moderated or removed, after which the cure may be accelerated by the substitution of some other remedy, such as one of the empyreumatic oils.

One of the most useful of the local applications in cases of lupus vulgaris is the *solid nitrate of silver*, if it is energetically applied. A pointed piece of caustic is made to penetrate the lupous deposit so as to destroy it completely, and to reach the healthy tissue beneath; after which no dressing is required, as the black crust which forms is sufficient protection from the air. When the crust falls or is removed, the caustic may be reapplied, and the treatment continued until all trace of lupous deposit has disappeared.

We are indebted to Mr. Higginbottom, of Nottingham, for the introduction of the use of a *concentrated solution of nitrate of silver* in the treatment of erysipelas. The affected surface is washed with soap and warm water, in case the skin should be oily; and afterwards with warm water, to remove all traces of soap (which would decompose the caustic), and dried with a soft towel. A solution of nitrate of silver, in the proportion of a scruple to a drachm of water, is painted over the affected skin and about an inch beyond the limits of the inflammation, and allowed to dry. If, on the

following day, any part is not blackened, thus showing that it has escaped the action of the caustic, the solution must be reapplied at that part; and if the inflammation has a tendency to invade new surfaces, these must in turn be attacked. If blisters form as the result of the caustic application, they need not be interfered with; but if they are the result of the erysipelatous inflammation, they should be opened, and the solution lightly applied to the abraded surface. It is scarcely necessary to add, that if the scalp is implicated the head should be shaved, as a preliminary to the subsequent treatment. Should the scalp be too sensitive to permit of this, the hair must be removed as thoroughly as possible with a pair of scissors. The extent of the inflammation upon the head can often only be detected by the occurrence of pain and pitting on pressure.


It sometimes happens that parts affected with lupus are attacked by erysipelas, and when the inflammation subsides the lupus is very much improved; and accordingly it has been recommended

by Hardy to apply a thin layer of a *strong ointment of red iodide of mercury*, in the proportion of equal parts of the mercurial and of lard, in cases of lupus vulgaris,—to excite an artificial erysipelas, in fact. This treatment, which may be repeated when the inflammation resulting from the previous application has subsided, is often followed by marked improvement,—but it is a painful process.

I have often derived the greatest benefit in cases of lupus from the use of the *acid nitrate of mercury*, which may be applied at intervals of from ten days to a fortnight. The crust which forms after the part is touched with the acid is sufficient protection to the ulcerated surface without any dressing. Under its influence the lupous deposit often vanishes, the profuse granulations disappear, and the ulceration heals.

Another method of treating cases of lupus vulgaris is by destroying the deposit by means of the *actual cautery*, with the aid of the instruments delineated in the accompanying woodcut, recommended to me by my friend Mr. Jonathan

Hutchinson. Patients who have been treated in this way do not seem to suffer nearly so much



Size $3/7$ ths,

as from the use of caustics; but on the whole I am inclined to think that the deposit is more

likely to recur than after the use of nitrate of silver.

In the treatment of obstinate circumscribed eruptions, *blisters* are often of great value; but then they must be applied *over* the affected part, not in the vicinity of it; that is to say, they are of no use as counter-irritants. Indeed, their application in the vicinity of an eruption sometimes leads to the development of a similar eruption on the blistered surface. This is especially true of the eczematous group of skin diseases.

The affected part may be readily blistered by painting it with a solution of a drachm of corrosive sublimate in an ounce of spirit: it is a painful process, however, and the danger of salivation is by no means slight; so that, if used at all, it should be restricted to very circumscribed, obstinate, syphilitic patches. For blistering purposes I generally employ Smith's "emplastrum cantharadinis liquidum," the directions for the use of which are indicated upon the bottle. The treatment of indolent ulcers by means of blisters, as first recommended by Mr. Syme, is so well known and

so thoroughly appreciated, that I need not dwell upon it. It is not so well known, however, that good results are not unfrequently obtained in cases of lupus erythematodes and lupus non-exedens, especially in the former ; but then the blistering requires to be frequently repeated. In cases of tinea decalvans I know no remedy which is so likely to prove successful as the long-continued use of blisters, while the same treatment is sometimes effectual, though not so generally indicated, in cases of tinea tonsurans (ringworm of the head).

The following case illustrates the value of this treatment where we have to deal with obstinate circumscribed patches of eczema :—A woman, well advanced in years, came to the Glasgow Dispensary for Skin Diseases on account of an eczematous eruption of old standing, which covered the palmar surface of each hand ; she had likewise a tendency to eczema of the leg. The eruption on the palms assumed that form which I have described under the name of eczema rimosum, the fissures being very numerous and deep, and the infiltration of

the skin considerable. She complained of itching and of pain; but the latter, on account of the fissures, predominated. Owing to the stiffness and pain, the hands were kept constantly in a half-closed position, and she was unable to use them. The effect of a single blister was astonishing. The eruption disappeared completely, and the patient returned with joy depicted in her countenance, not unmingled with pride, and opened and closed her hands with perfect facility.

Where the disease is obstinate, but too extensive to be blistered at one time—in cases of tinea decalvans implicating the whole of the head, for example,—the part may be attacked piecemeal, or an ointment containing cantharides may be used instead, as in the following prescription:—Powdered cantharides, two drachms; prepared lard, one ounce; red iodide of mercury ointment, one ounce. Melt a little, and paint the part as firmly as possible, short of causing pain or vesication, night and morning.

Iodine is chiefly valuable in the treatment of strumous affections of the skin. The use of the

tincture and ointment in cases of strumous enlargement of glands is too well known to need description, so that it will be enough to give one or two illustrations of the value of this remedy in other affections. In cases of lupus, especially of lupus erythematodes, the long-continued use of a concentrated solution of iodine and iodide of potassium may effect a cure. The following is the formula which I generally employ:—Iodine and iodide of potassium, of each half an ounce; glycerine (Price's), one drachm. Paint the part, and repeat daily till a thick red skin forms, or till the application becomes painful; then omit till the effect of the previous applications has passed off.

Again, in cases of tinea tonsurans (ringworm of the head) the following combination of iodine and tar, as recommended by Mr. Coster, has often a good effect, although in very strumous subjects it may lead to suppurative inflammation, and must then be abandoned: Pure iodine, two drachms; oil of tar (specific gravity .853), one ounce. The iodine and oil of tar should be gradually

and carefully mixed, otherwise a considerable amount of heat will be generated and the iodine dissipated.) "In the course of a week or ten days," writes Mr. Coster, "the scarfskin separates, and generally leaves a healthy surface beneath. I usually find that one application is sufficient to effect a cure where the disease is recent ; but if it be very chronic, or of several months' duration, it needs to be repeated perhaps three or four times."¹

There is a class of remedies denominated *parasiticides*, from the power which they are supposed to possess of destroying the animal and vegetable parasites which attack the skin. With regard to the first, there is a long list of remedies from which we may select. It is altogether out of the question, however, to enter fully into this subject ; so that one or two illustrations must suffice. These are selected either on account of their being well-known and well-tried remedies, or on account

¹ A Practical and Theoretical Treatise on the Diseases of the Skin. By George Nayler, F.R.C.S., p. 123. London : Churchill. 1866.

of their not being in common use, although equally valuable.

In cases of phtheiriasis (the disease due to the presence of lice), one of the best of all remedies, because one of the cleanliest, safest, and most effectual, is a lotion of corrosive sublimate, such as I have indicated in speaking of mercurial preparations. (See page 99.) When the lice attack the body (phtheiriasis corporis), I often prefer an ointment of staphisagria, of which the following is the formula :—Powder of staphisagria, one ounce ; lard, three ounces : digest for three hours, and strain.

In the treatment of scabies, as before remarked, the most universally employed remedy is sulphur, but one which may often be dispensed with ; for although it kills the acari, it irritates the skin, and thus tends to aggravate the artificial eruptions. I very much prefer, therefore, sponging the whole body night and morning with a solution of chlorate of lime, as recommended by Dr. Christison ; or anointing it every night for three successive nights with an ointment of styrax, of which the following is the formula :—Liquid

styrax, one ounce ; lard, two ounces: melt and strain. Or Schultze's modification of Pastau's prescription may be employed, and which is the following:—Liquid styrax, one ounce ; rectified spirit, two drachms ; olive oil, one drachm. This is enough for one patient. The styrax not only kills the acari as effectually as the sulphur, but also, instead of aggravating, soothes and favours the removal of the eruption called forth by the scratching.


There is much difference of opinion as to whether those local remedies which are used in cases of skin disease dependent upon fungous growths are really parasitocides, or do good merely in virtue of their stimulating action ; but, without offering any decided opinion on this point, I think it must be admitted that their stimulating qualities have something to do with the good result. Most of these have been already alluded to under other sections, such as sulphur, mercurial preparations, and empyreumatic oils and their derivatives. It may be well, however, to mention two others of proved utility, and more recently

introduced—namely, sulphurous acid and the hyposulphite of soda. The first may be used pure, or diluted with water; the second dissolved in water in the proportion of from half a drachm to a drachm to the ounce. The affected parts, after the removal of epithelial *débris*, &c., should be sponged with these two or three times a day.

It must never be forgotten that when vegetable parasitic skin diseases implicate hairy parts, and when the hairs are attacked, we are unable to reach the fungous matter which lies imbedded in their roots, and, accordingly, epilation must frequently be superadded. Indeed, as far as my experience goes, it is quite essential in the treatment of tinea sycosis (ringworm of the beard), and of tinea favosa when the hairy parts are attacked.

The woodcut on the next page illustrates the kind of instrument which should be used for the purpose. The forceps should be about three inches long, and should be made so that the two extremities come together very exactly, and do not slide the one upon the other. Each extremity should be furnished on the inside with denticula-

tions, after the manner of a file, and these should be very fine, else the brittle hair may be cut across by them. The forceps should not have a strong



spring, otherwise the hand soon becomes fatigued in using them.

It is necessary to avoid extracting the hairs except in the direction of their axes, else they are very apt to break; and not more than half a dozen hairs at the most should be seized at one time, and even these must be in close proximity to one another. It is advisable, too, to clean the points of the forceps often, for the *débris* which accumulates between the denticulations is apt to cause the forceps to slip when the hair is grasped. The pain of epilation may be diminished by rubbing oil of cade into the parts night and morn-

ing, as suggested by Bazin (which, at the same time, acts the part of a parasiticide); but, for

my part, I prefer to keep them constantly lubricated with almond or cod-liver oil, which not only diminishes the pain, but also causes the hair to grow more healthily and renders it less likely to break. After one complete epilation, the treatment should be stopped for a week or so, after which partial epilations may be required, at those parts, namely, where the disease reappears owing to the roots not having been completely removed.

It seems unnecessary, as recommended by some, to destroy the clothing of a patient affected with a parasitic skin disease, although it may sometimes be the most convenient way of removing sources of reinfection,—in the case of the lining of bonnets, for example, when patients are labouring under ringworm of the head. But if we wish to ensure the non-recurrence of the disease—in cases of scabies, for instance—it is necessary to disinfect the clothing, which can be done in a variety of ways, such as steeping it in boiling water, or exposing it to the fumes of sulphur, or to the influence of very hot air.

CHAPTER V.

B.—CONSTITUTIONAL TREATMENT.

BEFORE entering into details with regard to the constitutional treatment of diseases of the skin, a few preliminary observations are desirable. And in the first place, let me remark that a disease which is purely local cannot be benefited by any internal medicine, except arsenic, which acts specially upon the skin, while local treatment is almost always preferable to it. On the other hand, if an eruption is symmetrical—thus indicating, as a rule, its constitutional origin—or diffused over an extensive surface, it is much more likely to yield to constitutional treatment. Again, if an eruption is coming out rapidly, even when there is no constitutional reaction, it generally requires what is denominated a cooling

treatment, and is likely to be aggravated by the use, at the outset, of iron, arsenic, and such remedies. It must also be borne in mind that if any derangement of internal organs, of the kidneys, digestive organs, &c., complicates a skin affection, remedies specially directed against the skin disease—tonics, for example—are almost certain to aggravate it.

As long as an eruption continues decidedly to improve, and if the general health is not suffering from the use of the remedy, no radical change should be made in the treatment; but if, as sometimes happens, the patient tires of the long continuance of one prescription, the remedy may be given in another form. Further, if a particular medicine is decidedly indicated, and if the patient starts objections to it without good reason, and if there is reason to believe that no other medicine has any chance of being equally effectual, it is sometimes allowable to give it in a concealed form. (It is in order that the patient may not know that he is taking opium that the name of the opium pill of the Pharmacopœia has been

changed from "pilula opii" to "pilula saponis composita.")

In conclusion, let me say that the most judicious treatment is very apt to prove abortive if the diet and regimen are not carefully attended to, in regulating which we must be guided by broad general principles. And let me add—and this is a point which is too often ignored—that sea-air and sea-water are very apt to prove prejudicial in the subjects of skin affections, with the exception, perhaps, of those labouring under strumous diseases.

Purgatives or aperients are of service in a large proportion of cases, at the outset at all events; and many can be cured by the exclusive use of them. This is especially true of the eczematous group when seen in the early or acute stage, and when associated with digestive derangement or constipation. In doubtful cases it is safer to inaugurate the treatment with aperients; but it must never be forgotten that where debility is well marked, or where there is a decided tendency to ulceration, great caution

is required in the use of purgatives. In the employment and selection of purgatives and aperients we must be guided by general principles.

A very excellent aperient is the well-known sulphate of magnesia and sulphuric acid mixture ; to which, if there is a gouty tendency, a little wine of colchicum, or, if anæmia, sulphate of iron, may be added as follows :—Sulphate of magnesia, three ounces ; dilute sulphuric acid, an ounce and a half ; sulphate of iron, three drachms ; simple syrup, four ounces ; tincture of ginger, one ounce ; infusion of quassia to twenty-four ounces : a tablespoonful, in a good deal of water, three times a day.

Another very useful tonic aperient, especially if the bowels are easily moved and there are indications of nervous debility, is the following :—Phosphate of soda, three ounces ; dilute phosphoric acid, two ounces ; syrup of ginger, four ounces ; compound infusion of gentian, eight ounces ; distilled water to twenty-four ounces : shake the bottle : a tablespoonful, in a large wineglassful of water, to be taken three times a day.

In each case the dose must be so regulated that

the bowels are kept freely open. If purging is induced, it must be reduced ; while, if the bowels are not acted upon, a pinch of sulphate of magnesia may be added to each dose in the case of the first prescription, a little phosphate of soda in the second.

In acute cases, one or two full doses of calomel, followed by castor-oil or a seidlitz powder, is often beneficial at the outset ; but prolonged purgation must in every case be avoided, for although it may remove the eruption for a time, it returns whenever the treatment is stopped, and, by weakening the patient, may indefinitely prolong the disease.

We are told that sulphur is the great blood depurant in the case of diseases of the skin ; but for my part, I am as little partial to it when administered internally as when used as a local application. When it does good, it is generally in virtue of its purgative action ; while it has this drawback, that it is converted into sulphuretted hydrogen, so that the secretions have a disagreeable odour. It is usually given in electuary with

acid tartrate of potash, but it is better to prescribe one of the natural mineral waters containing it; and the fact that some of these do not act as purgatives must not be taken to disprove my assertion with regard to the *modus operandi* of sulphur, for the benefit which accrues may be due to the combination of salts held in solution. Those of Harrogate and Moffat in this country and of Aix-la-Chapelle, Enghien, Bareges, and Luchon on the Continent, have the greatest reputation in this respect; and while some of these waters may be had from the chemist, it is always more judicious, when it can be effected, to send the patient to the spring itself, for he is thus certain to get the waters fresh and pure, and, away from home and the fatigues and anxieties of business, his body is at the same time invigorated and his mind refreshed.

Diuretics are indicated if the kidneys are torpid, especially in the treatment of the erythematous and eczematous group and in acute cases. Their diuretic action, however, does not always account for the good result observed: thus alkaline diu-

retics, such as the bicarbonate of potash, or neutral (which in the system are converted into alkaline) salts, such as the acétate of potash, probably do more good in virtue of their alkaline reaction. And this leads me to remark that *alkaline medicines* are especially useful in the treatment of skin diseases occurring in rheumatic¹ and gouty subjects, and in persons who are martyrs to acidity, or in whom there is a tendency to the deposit of uric acid and of urates in the urine. The alkaline and neutral preparations which are most used are salts of potash, such as the acetate, bicarbonate, and citrate; but if a stimulant is required, I generally prefer the carbonate of ammonia; and it will often be found of advantage to combine these remedies with arsenic, or, if there is a gouty tendency, with colchicum. The following is a useful combination:—Carbonate of ammonia, one ounce; solution of arsenic, three drachms; syrup of ginger, six ounces; infusion of cascarilla to

¹ A somewhat extensive trial of alkalies in the treatment of rheumatic fever, and especially in lingering cases, in which the effects of remedies can be more readily appreciated, has convinced me that they are often of service.

twenty-four ounces : a tablespoonful in a third of a tumblerful of water three times a day.

It must never be forgotten that alkalies should be given largely diluted, and also some time before food ; for although the administration of a small quantity of an alkali seems to cause an increased flow of gastric juice, a large quantity may neutralize its acidity. The dose should be so regulated as to keep the urine constantly alkaline as tested with litmus paper ; and the medicine should be steadily continued, unless it disagrees, until some time after the disease has disappeared.

Sedatives and narcotics are indicated under the same circumstances as in the treatment of disease of other organs. But it is necessary to bear in mind—and the remark applies not merely to skin diseases—that sleeplessness is often the result of debility, in which case tonics, especially iron and arsenic, are the best narcotics. Sedatives and narcotics are of no use for the relief of irritation of the skin, except in so far as they may induce sleep ; indeed, if they derange the stomach they are very apt to increase the irritation.

But, as is well known, opium in small doses, frequently repeated, is of value in the treatment of affections of the skin occurring in broken-down subjects, especially in cases of ulceration; but then, they do good in virtue of their stimulating, rather than of their narcotic, properties.

In genuine cases of urticaria perstans, that is, nettle-rash, recurring from day to day and from week to week, when it is independent of local causes, and when no constitutional derangement (disorder of the digestive organs or the like) is present, sedatives are sometimes curative, especially bromide of potassium in full doses.

Lastly, it need only be mentioned that the neuralgia which so frequently accompanies and follows attacks of zona (shingles) is best relieved by the subcutaneous injection of morphia.

Cod-liver oil is indicated in the treatment of strumous diseases of the skin, although it is of more value in preventing fresh outbreaks than in removing existing manifestations. The following case illustrates the above remark:—

A little boy was brought to the Glasgow Skin

Dispensary who had an elevated, warty, strumous patch about two and a half inches long by one and a half broad on the buttock—that disease which I described recently¹ under the name of lupus verrucosus. No local treatment was prescribed, but he was ordered to take a tablespoonful of cod-liver oil three times a day. The eruption improved very slowly, and it was about nine months before it had completely disappeared.

Cod-liver oil is also indicated in the treatment of skin diseases, although they are not strumous, provided they occur in strumous, half-starved, or broken-down subjects.

In dispensary practice we are often consulted by persons who have long been out of employment, whose diet has been very defective, and whose bodies are infested with lice. The use of cod-liver oil in full doses, without the use of parasitocides, often removes the disease by supplying the place of a nourishing diet, and by improving the general health, *i.e.* by rendering the soil—uncongenial to these unsavoury companions.

¹ Journal of Cutaneous Medicine, vol. i.

Two more illustrations of the beneficial effects of cod-liver oil may be given.

A child, fifteen months old, was brought by his mother to the Glasgow Dispensary for Skin Diseases on Oct. 9th, 1862, affected with eczema impetiginodes. The eruption implicated almost the whole of the body, was very itchy, constantly exuding, and studded with crusts. The child was dreadfully emaciated, "just skin and bone," as the mother remarked. It could neither sleep nor eat, and was so weak that it had to be brought upon a pillow. The case looked hopeless, and, indeed, the child had been given up by the previous attendant; but, acting upon what I have observed in similar cases, twenty drops of syrup of the iodide of iron in a teaspoonful of cod-liver oil were prescribed, to be repeated thrice daily, and the dose of the oil to be gradually increased to a tablespoonful. "On Oct. 16th the child was better. The skin being still itchy, however, a lotion of dilute hydrocyanic acid, containing twenty minims to the ounce of water, was ordered, to be used thrice daily as a palliative. The oil

was omitted for a week, as it produced purging. With this exception, the oil and iron were steadily continued till November 17th, about five weeks after the commencement of the treatment, when the mother brought the child, out of gratitude, to show how well it was. There was hardly a vestige of the previous eruption, with the exception of a few dry crusts and discoloured spots on the buttocks, which were rapidly disappearing. The child appeared to be in robust health; it was quite plump, and its cheeks rosy; its skin soft and white; its appetite very good; and its sleep sound and refreshing. The medicine was to be continued for another month."

It may be well to remark in this place that if cod-liver oil is taken greedily by infants, it is almost certain to do good, although the reverse does not hold.

A man, aged about fifty, came to the Glasgow Dispensary for Skin Diseases in the year 1864, on account of an attack of psoriasis. He had a severe posterior curvature of the spine in the dorsal region; and, being out of employment,

his diet was of a very inferior description. The eruption was very extensive, and had existed uninterruptedly to a varying extent for twenty-eight years; and yet, in the space of six weeks, under the influence of full doses of cod-liver oil alone, it had all but entirely disappeared; and I then lost sight of him.

It will thus be seen that I differ from Hebra, who limits the value of this medicine to the treatment of strumous diseases; and from many others who, on the other hand, laud it in the treatment of an immense number of cases of skin disease.

It is unnecessary to dwell upon the use of *iodine and its compounds*, as these are indicated in the same class of cases as cod-liver oil. Of course this remark does not apply to the iodide of potassium, to which reference will be subsequently made.

CHAPTER VI.

AS so many affections of the skin occur in connection with, if not in consequence of, the system being below par, *tonics*, especially iron and arsenic, are frequently indicated. With the exception, however, of *arsenic*, which is the great cutaneous tonic, it is unnecessary to point out the indications for their use, as these are the same as in the treatment of disease in general. There is much difference of opinion as to the use of arsenic; and while I am altogether opposed to those who are inclined to disparage its virtues, I am equally opposed to the view that almost all affections of the skin should be treated by means of it. At the present time I must content myself with a very few remarks, which, for the sake of brevity, I make in the form of aphorisms :—

1. Arsenic, judiciously administered, is as safe a medicine as any in the Pharmacopœia, and may often be used for months without injury to the general health.

2. It often requires to be continued for many weeks, and sometimes the disease seems to resist its action for a considerable time, when all of a sudden improvement occurs, followed by a rapid cure.

3. It requires to be given in proportionately larger doses to children than to adults.

4. Infants may be subjected to its influence by administering it to their nurses.

5. The dose should be at first small, and not increased, as a rule, for some time. Then it may be gradually increased till the medicine disagrees, or till the disease begins to yield, when it may as gradually be diminished.

6. It should not be omitted altogether without very good reason, but may be tried in smaller doses or in another form, or omitted for a few days, till the bad effects have passed off.

7. Puffiness of the face, or irritation of the eyes, or such-like physiological effects, if slight in

degree, should not lead us to discontinue the medicine; indeed, it is sometimes only then that its beneficial action on the disease is observed.

8. It is decidedly contra-indicated in acute cases; and when its use is followed by marked increase of the irritation of the skin (itching, heat, &c.), the disease is probably not in a state to be benefited by it.

9. It is generally more rapidly effectual if the disease, though in a chronic state, is recent; and the first attacks yield more readily to it than subsequent ones, as a rule.

10. It is contra-indicated in most cases which are complicated with digestive derangement.

11. It is apt to produce bronchial catarrh, so that patients should be warned to avoid exposure to cold while taking it; and for this reason it is generally contra-indicated in persons labouring under bronchitis.

12. In exceptional cases it may be given with benefit in large doses, as in the following cases:—

On May 11th, 1861, I was sent for to the country for the purpose of seeing a little girl,

aged about ten, who had been suffering for about three months from a papulated eczematous eruption, principally affecting the back. When I saw her she was confined to the sofa, and at that time her whole back, from the neck to the hips, presented an enormous ulcerated surface. The ulceration was quite superficial, and presented a slightly papulated aspect. It had all the appearance of a superficial ulcer from a burn which was gradually contracting and healing at the edges. At the margin, also, papules and vesicles, containing opaque serum, were detected. Papules were likewise scattered thinly over the body, but especially on the brow. From the surface of the sore semipurulent matter was exuding. The little girl had been able to run about till within a week of the above date, since which time she had been confined to the sofa. Her general health was, however, good, except that she had suffered a little from the confinement and from the irritation of the sore.

Dr. Robert Stewart, of Coatbridge, saw the patient with me, and we agreed that the sore

should be dressed with cod-liver oil, and Fowler's solution administered in gradually increasing doses.

I am indebted to Dr. Stewart for acquainting me with the result of the treatment. In a letter dated October 22nd, 1861, he wrote:—"After you saw her she commenced with two drops of Fowler's solution three times a day. Each dose was increased by a drop each day, so that latterly she was taking thirteen drops of Fowler's solution three times a day, which had the most charming effect, and produced a decided cure. Altogether she must have taken, in the course of six or seven weeks, two and a half ounces of the solution. I saw her regularly, and there never was a bad symptom."

I have at present under my care, in the Glasgow Royal Infirmary, a little girl, aged nine, who has been affected with pemphigus in a most aggravated form for two years, and who has been taking ten minims of Fowler's solution thrice daily for many weeks with perfect impunity, and with the effect of entirely removing the bullous eruption. It is

exceedingly probable, however, that such large doses would induce catarrh of the mucous membrane of the respiratory tract if the patients were not kept within doors and protected from draughts.

13. It sometimes requires to be given during meals, or immediately after food is taken, for when administered on an empty stomach it occasionally deranges the digestive organs; and it is often better tolerated if given along with a bitter infusion.

14. It should not, as a rule, be entirely discontinued until some weeks have elapsed since the complete disappearance of the eruption.

15. There are few chronic diseases of the skin of constitutional origin—provided they are not syphilitic—which may not be benefited by it (although often other treatment is to be preferred to it), but it is especially valuable in psoriasis, pemphigus, lichen ruber, pityriasis rubra, and in many cases of eczema; unless contra-indicated as above.

The preparations which I am most in the habit of using are liquor arsenicalis (Fowler's solution), of which the medium dose for children is two, for

adults five, minims thrice daily, and Asiatic pills, of which the following is a modified formula:— White arsenic, two grains; black pepper and glycyrrhiza powder, of each half a drachm; with a sufficient quantity of mucilage: divide into thirty-two pills, one to be taken two or three times a day.

Empyreumatic oils and their derivatives.—In former days tar-water and tar pills appear to have been favourite remedies in the treatment of chronic diseases of the skin, but these, as is well known, have long ago fallen almost entirely into disrepute. Being struck, however, with the powerful influence of tarry preparations in the treatment of certain chronic affections when applied to the skin, and feeling convinced that their action is not a mere local one; being sure, too, that when a remedy has been at any time in favour with the profession there must be some foundation for its use, I determined to give them a trial.

The preparations which I have employed are *pix mineralis* (coal-tar) and *pix liquida* (wood-tar); and of these the latter is less likely to

derange the digestive organs. The medicine should be given in treacle, or in the form of capsules; but on no account in the form of pills, as they are apt to become hard and consequently useless.¹ The dose is two or three minims thrice daily, and it may be gradually increased to twenty if necessary. If the digestive organs are in good order and the bowels free, and if too large a dose is not given at the outset, the medicine agrees perfectly. It is useful in cases of chronic psoriasis implicating a large extent of surface; also in chronic eczema, especially if associated with chronic catarrh of the stomach or bowels, or with chronic bronchitis—morbid conditions which it sometimes modifies or removes entirely.

A young lady, aged about twenty-six, of good constitution, consulted me on the 15th July, 1867, on account of an attack of the so-called acne rosacea, of many months' duration, and implicating the whole of the face, which was of a deep crimson tint; she had also a few spots of psoriasis upon

¹ The capsules are prepared for me by Mr. Robert T. Dun, 288, Argyle Street, Glasgow.

her brow and one upon her knee. An arsenical course had been prescribed for her two years before, on account of the psoriasis, by Mr. Erasmus Wilson, and she had continued to take it ever since on her own responsibility, for she found that whenever she stopped it the psoriasis reappeared. The arsenic was now discontinued; a tonic aperient mixture was given, and an ointment of rumex and sulphur (*ante*, p. 52) prescribed for the face. On the 30th August the psoriasis was very decidedly on the increase, and she desired to resume the arsenic. Instead of this, however, she was ordered to take *pix liquida* in capsules; at first in three-, gradually increased to fifteen-minim doses, thrice daily. On Oct. 8th, when I next saw her, the tendency to the aggravation of the psoriasis had been completely checked; and on the 10th December the eruption had almost disappeared, while the face had resumed its normal appearance. The cure of the *acne rosacea* was no doubt due in part to the use of the ointment; but the psoriasis was checked by the *pix liquida*, as no outward treatment was resorted to for it. Moreover, she found that when-

ever she stopped the tar, just as in the case of the arsenic, the psoriasis tended to return; and, seeing that her health was perfect in every respect, and that she suffered no inconvenience from the use of the capsules, I allowed her to continue them, and, for all I know, she may be doing so yet.

In this case the *pix liquida*, like the arsenic, only kept the disease in check; but then it was a very chronic and exceedingly inveterate one. Did time permit, I could relate other cases in which *pix liquida* was more speedily and more permanently effectual; but I have selected this one, as it appears to me to be very instructive.

Within the last two years, encouraged by the results of the tar treatment, I have tried, on an extensive scale, the internal administration of *carbolic acid*, which, as might be expected, is indicated in the same class of cases as the tar preparations. I prescribe it in doses of from three to ten grains thrice daily, and usually in the following form:—Crystallized carbolic acid, three drachms; glycerine, one ounce; distilled water to six ounces: a teaspoonful in a *large*

wineglassful of water thrice daily, some time before food.

If the same precautions are observed as in the case of the tar, the remedy rarely disagrees, although transient giddiness is sometimes complained of a minute or two after it is taken, especially after the morning dose. If no improvement takes place within two or three weeks, it generally does not succeed, or at all events not without increasing the dose; but a slight aggravation of the eruption for a few days after the medicine is commenced is by no means an unfavourable symptom.

Carbolic acid is especially useful in cases of chronic psoriasis when the patches are not much infiltrated, and not unfrequently I have succeeded in removing the eruption after a fair trial of arsenic, although, on the whole, it is not so useful as the latter. Those, however, who have much to do with the treatment of psoriasis cannot but consider it a great boon that another remedy has been added to the very limited list of internal medicines which are capable of modifying that most obstinate disease.

The following cases—and I could quote many others—may prove interesting in connection with the preceding remarks.

CASE I. *Psoriasis*.—Mary G——, aged nine, school-girl, was admitted to the Glasgow Skin Dispensary on January 4th, 1869. Her mother stated that before the appearance of the eruption, which occurred six weeks previous to her visit, she was not at all in good health, but that immediately thereafter she recovered completely.

Appearances on admission.—The left leg was thickly covered with spots, which were, for the most part, the size and shape of a florin. The left thigh also was the seat of numerous spots, varying in size from that of a pea to that of a shilling. The nates were studded with them, and these for the most part were the size of peas. The extensor surfaces of the arms were covered with spots about the size of a sixpence, whilst very few were found on the flexor surfaces. A few small spots were seen on the right cheek. All the patches were circular in form; the larger ones presented a pinkish tint, while the smaller were covered with

typical silvery scales. The eruption was extending rapidly, but was not in the least degree itchy. Ordered the following mixture :—Crystallized carbolic acid, one drachm and a half; glycerine, a sufficient quantity; distilled water to six ounces: a teaspoonful in a wineglassful of water three times a day, on an empty stomach.

The following reports of the subsequent progress of the case were taken :—

January 18th, 1869.—For a few days after the medicine was commenced the eruption “hacked” more, was stiffer and more painful. Since then it has become less red, and the scales are falling off very rapidly. Medicine agrees, “and she is eating much more.”

February 1st.—Continue, as medicine agrees. Eruption becoming paler and less scaly, and no new spots.

18th.—Mother says she never improved so much as she has done the last fortnight. Very little to be seen now, except the stains of previous eruption.

This patient did not return.

CASES 2 AND 3. *Psoriasis*.—Agnes L——, aged fifteen, factory girl, was admitted to the Glasgow Skin Dispensary on December 14th, 1868. The first outbreak, which commenced immediately after an attack of scarlet fever six years previous to admission, covered almost the whole body, and got well under medical advice. The attack on account of which she came to the Dispensary was of four weeks' duration, and commenced upon the elbows.

Appearances on admission.—On the right elbow there was a round patch about the size of a florin, covered with silvery scales; while numerous small spots of the same character, and varying in size from a pin's head to that of a pea, were scattered over the fore-arm. The left arm was similarly affected, but the patch on the left elbow was much larger, being about three inches long and half an inch broad. On the right knee was a patch about the size of half-a-crown, and numerous spots about the size of a split pea were scattered over the leg, all of which were covered with silvery scales. The same appearances were observed on

left knee and leg. The eruption was increasing rapidly, and was very itchy. Her general health was good. Ordered, seventy-two grains of crystallized carbolic acid in twenty-four ounces of distilled water: a tablespoonful in a wineglassful of water three times a day, on an empty stomach.

As her sister Ann, aged seventeen, was reported to have the same disease and to be otherwise in good health, the same medicine was prescribed for her, and a message was sent to her to come to the Dispensary the following week.

The following were the subsequent reports:—

December 28th, 1868.—In both the above cases the spots came out more rapidly for the first week, but none since; and the old spots and patches are beginning to disappear.

January 11th, 1869.—Improvement very marked, especially in Agnes.

February 8th.—Very great improvement in the case of Agnes since last visit. Decided improvement in the case of Ann, although not so marked; therefore let Ann increase the dose to a tablespoonful and a half.

March 17th.—Agnes all but cured. Ann very greatly better, and disease progressing rapidly towards a cure under the increased dose.

These patients did not return. (Reported by Mr. J. D. Walker.)

CASE 4. *Psoriasis.*—A lady aged about thirty-five, who had undergone much anxiety and fatigue in nursing a sick child, was attacked for the first time by psoriasis. The patches, on an average about the size of florins, were of a dusky red tint, but not scaly. They were very numerous, especially upon the arms and legs. She felt weakly, and was incapable of much exertion, and the feeling of fatigue and lassitude was very marked at times; otherwise her health was good. She was subjected to an arsenical course by her medical attendant, but without benefit. She then consulted me, when, the arsenical course having failed, four grains of carbolic acid in solution was prescribed thrice daily. There was immediate improvement. The feeling of lassitude and exhaustion diminished, the eruption began to fade, and in a few weeks she was quite well.

CHAPTER VII.

IN these days of scepticism, when we hear so much of the successful treatment of syphilis without mercury, it becomes those whose experience justifies them in giving an opinion, to do so clearly and without reservation. It is my firm conviction, then, that while mercury is of little use in the treatment of non-specific affections of the skin, except in so far as it is of value as a purgative or corrector of digestive derangement, it is invaluable, in appropriate cases, in syphilitic affections; and further, that no one who has not had ample opportunities of testing its efficacy in syphilitic affections of the surface, where we have the eye to guide us, is qualified to form a correct opinion as to its value in syphilis.

We are justified in bringing the patient fairly under the influence of mercury under the three following conditions, although it is not always necessary to touch the gums, and certainly in no case should salivation be courted.

1st. When we have to treat one of the earlier of the cutaneous manifestations of syphilis in a person of sound constitution.

2nd. When the syphilitic poison has not only attacked the skin, but also some delicate organ, such as the eye.

3rd. When, in a person of sound constitution, we have to deal with a circumscribed syphilitic eruption of old standing, which resists external applications and iodide of potassium in full doses.

A gentleman, resident in the West Highlands, consulted me a couple of years ago with regard to an obstinate syphilitic eruption of old standing, implicating, and limited to, the soles of his feet. The patches were slightly coppery in tint; their edges were composed of segments of circles; and the skin was for the most part thick and

almost horny. He had taken iodide of potassium *ad nauseam*, and had tried all kinds of external applications, including the use of blisters, under the direction of one of the most celebrated surgeons of the day; but to no effect. He was an exceedingly strong and healthy man, and I had therefore no hesitation in giving him a course of mercury. Within three weeks the eruption had completely disappeared, and there has been no return since, while his general health remains as perfect as before.

Instead of administering mercury by the mouth, the process of subcutaneous injection, as recommended by Lewin and others, may be tried—a mode of administration from which I have obtained excellent results. Four grains of the perchloride of mercury should be dissolved in an ounce of distilled water, and of this from seven and a half minims at the least (*i.e.* a sixteenth of a grain) to thirty at the most (*i.e.* a quarter of a grain) should be injected once daily. It is indifferent where the injection is made, except that the least sensitive parts and

the parts least liable to pressure should be selected. It possesses the following advantages:—

1. The dose can be regulated with the utmost precision. 2. The medicine has no tendency to derange the digestive organs. 3. It acts with greater rapidity than when the perchloride is administered by the mouth. 4. A much smaller quantity is sufficient to produce the desired effect. 5. It does not usually produce salivation, although stomatitis is a frequent result. The process has, however, the following disadvantages:—1. The operation is somewhat painful, and much pain is experienced for several hours after each injection; so that patients sometimes object to its continuance. This, however, may be counteracted by adding one-sixth of a grain of morphia to each injection. 2. In exceptional cases—once in about two hundred injections (Lewin)—the operation is followed by circumscribed inflammation and abscess.

Another mode of employing mercury is in the form of the *mercurial vapour bath*, which is specially valuable in the treatment of syphilitic

ulcerations of the skin, and also of syphilitic eruptions occurring in cachectic subjects; but I am not prepared to coincide with those who laud it as the most effectual, although it is undoubtedly the safest, method of bringing the system under the influence of mercury in every case. The preparations which are most frequently employed for this purpose are the subchloride and the bisulphuret; of which half a drachm of the former and two drachms of the latter may be used for each bath. The mode of carrying out this treatment, and its special advantages, have been fully discussed by Mr. Langston Parker and Mr. Henry Lee, to whose writings I must refer the reader for further details.

In syphilitic eruptions occurring in infants, the subjects of hereditary taint, mercurial treatment is certainly curative, provided it is commenced before the poison has had time to produce profound syphilitic cachexia. Indeed, there is no other treatment which appears to have any material influence over the disease; so that, in my

opinion, to withhold mercury in such cases is altogether unjustifiable.

As a rule, the most simple and the most satisfactory way of treating infants is to rub into the eruption, night and morning, an ointment containing mercury, such as I have indicated in a previous chapter, or to make use of the mercurial belt. A piece of mercurial ointment the size of a bean should be rubbed daily upon the inside of a piece of flannel two inches broad, and sufficiently long to surround the body of the child about the level of the umbilicus, and this should be worn until two or three weeks after the complete disappearance of the symptoms. Should the ointment, however, irritate the skin, and tend to produce an erythematous or eczematous rash (the so-called *eczema mercuriale*), the belt may require to be removed for a time, during which the ointment may be rubbed into some other part of the body.

Everyone is familiar with the cutaneous manifestations of infantile syphilis; but it does not seem to be generally known—at least, there is

no allusion to it in most dermatological works—that eruptions occurring after the infantile period in children and young persons are sometimes dependent upon hereditary syphilis. The diagnosis is often difficult, but the beneficial effects of antisyphilitic treatment sometimes serve to convert a suspicion of syphilitic taint into a certainty. Cases illustrative of what has just been stated have been referred to in a previous chapter (see Part I. chapter ii. pp. 35—40).

CHAPTER VIII.

WITH regard to the employment of *iodide of potassium* in the treatment of syphilitic skin diseases, I must content myself by laying down the following rules:—

1. The longer the interval which has elapsed between the contraction of the syphilitic taint and the development of the eruption, the more confidently may we substitute it for mercury.

2. If the patient is cachectic, it is, as a rule, to be preferred to mercury, except in recent cases of syphilis, when the mercurial vapour bath, or some such treatment, is more likely to prove successful.

3. The more extensive the tertiary eruption, the more certain it is to yield to the iodide of potassium; although to this rule there are numerous exceptions.

4. If there is any tendency to syphilitic disease of the nostrils or neighbouring parts, iodide of potassium should be withheld, or given with great caution, for, if it produces coryza, it is very apt to aggravate the morbid condition of the parts.

5. It should be given in full doses.

The last rule is one of great importance, and all the more so, seeing that one of the most distinguished surgeons of the present day recently recommended, in one of the medical journals, the administration of doses of three grains; while my own experience has led me to conclude that ten grains is the proper dose in the majority of cases, and that occasionally as much as thirty or forty, thrice daily, may be requisite. It is generally advisable to prescribe it in combination with a bitter, and in cachectic subjects a little iron is a valuable addition, as in the following prescription:—Ammonio-citrate of iron, three drachms; iodide of potassium, one ounce; syrup of ginger, six ounces, compound infusion of gentian, eight ounces; water to twenty-four ounces: a table-spoonful in a large wineglassful of water thrice daily.

The Alkaline Sulphites and Hyposulphites.—Some years ago Polli of Milan made experiments with the sulphites and hyposulphites of potash, soda, and magnesia. Having proved, by experiments—"first, that sulphites, when administered to a living animal, are carried, *as such*, into the circulation, and diffused all over the organism without the slightest inconvenience to the animal; and, secondly, that the presence of these salts in the liquids and solids of the body retards the putrefactive fermentation for a very considerable period he took two dogs of about the same size, and equally in good health; he fed them exactly alike for five days, with the exception of administering to one of them two grammes of sulphite of soda daily—the other dog getting exactly the same food, minus the sulphite. At the end of five days he injected into the femoral veins of both animals one gramme of pus taken from a fetid abscess occurring in a broken-down constitution. The operation in both cases was carefully performed, and the animals suffered but little. Immediately after the injection both dogs appeared

stupefied; they lay down and refused all food, remaining quite prostrate for twenty-four hours. On the following day, however, they both seemed a little better, and took some food. A second injection of pus was now practised on both animals to the same amount, but the first dog had meanwhile been getting two grammes of sulphite daily, while the latter was only getting plain food. The effect of the second injection was most interesting: both dogs were affected instantly alike; both were seized with stupor; in both the pulse was rapid and feeble, while the respiration was greatly accelerated; both dogs refused to eat; both lay down in a state of stupor; and, when made to rise and walk, they tottered and reeled across the room. The first dog, however, continued to receive, daily, a dose of two grammes of sulphite of soda, and in four days was so far recovered as to be able to eat his food with relish, while the wound in the femoral vein was rapidly healing. The other dog fared differently: he got no sulphites, either before or after the operation, and the result was that he daily became worse; the wound in the thigh

became gangrenous, the limb swelled up, and ten days after the second injection the dog died, with all the symptoms of typhus; the first dog being already about and well.”¹

One other experiment may be mentioned. Polli “took two large dogs, as similar as possible both in size and health, and, having administered to one of them eight grammes daily of sulphite of soda, he injected into the femoral veins of both dogs three grammes each of the muco-purulent discharge obtained from the nares of the same glandered horse which had served for a previous experiment. The first dog, which had received the sulphites, seemed at first to suffer the most from the injection. It at once fell to the ground as stunned, and its breathing was rapid and panting; but in a few hours it began to recover, and the following day it was able to eat. The second dog bore the operation better, and did not appear to sustain so severe a shock; but on the following day it began to mope, towards evening it was very

¹ Dublin Quarterly Journal of Medical Science, vol. xxxvi. p. 470 : article by Dr. de Ricci.

drowsy, and with difficulty it could be got to stand ; by the third day the animal's extremities had become œdematous and painful ; by the fourth, a purulent discharge was running from its nose and eyes, and the wound in the thigh was now almost gangrenous. On the fourth day the animal died, worn out by pain, fetid suppuration, and diarrhœa. The first dog was by this time completely recovered."¹

For further details I must refer those who are specially interested in this subject to Dr. de Ricci's paper, and to Professor Polli's work "On the Use of Alkaline Sulphites in the Treatment of Diseases depending on Morbific Ferments."

Acting upon these data, and believing that impurity of the blood lies at the root of those cases of recurrent furunculi in which no local or special constitutional cause can be detected, I made a trial of the hyposulphite of soda in this complaint, and in some instances I was surprised at the result, as in the following case :—

¹ Dublin Quarterly Journal of Medical Science, vol. xxxvi. p. 407 : article by Dr. de Ricci.

“A young man, of good constitution, but who for some months had suffered from a series of boils, appearing principally upon his legs, came for advice to the Dispensary for Skin Diseases. The complaint had not moderated in the least when I saw him; for the furunculi were coming out in rapid succession. I prescribed for him half a drachm of the hyposulphite of soda, thrice daily, in water; and from that day he never had another boil.”

Before giving the medicine we must make inquiry as to the condition of the general health, especially as to the state of the bowels, which are frequently constipated, and correct any derangement which may be present. It should be given largely diluted, as in the following prescription:—
Hyposulphite of soda, an ounce and a half; simple syrup, one ounce; cinnamon water to twelve ounces: a tablespoonful, in a large wineglassful of water, thrice daily, on an empty stomach.

Dr. de Ricci prefers the sulphite of magnesia for internal administration; “because it is not so unpalatable, and is less likely to produce diarrhœa

and because, in consequence of the atomic weight of magnesia, it contains, bulk for bulk, more acid than the soda salt." He cites a case of chronic pemphigus in an old gentleman over eighty, in which the external use of a saturated solution of sulphite of soda, and the internal administration of the sulphite of magnesia, had a remarkably beneficial effect.¹

It appears, then, that the alkaline sulphites and hyposulphites are of value in the treatment of some diseases of the skin; and that they are deserving of a more extended trial than has hitherto been accorded to them.

¹ Dublin Quarterly Journal of Medical Science, vol. xlii. p. 363.

INDEX.

A.

ABSORBENT powders, 85
 Acetate of potash, 136
 Achorion Schönleini, 27
 Acid nitrate of mercury, 118
 Acne vulgaris, 7, 9, 50
 treatment of, 98-103
 Actual cautery, 118
 Acute inflammation of skin,
 characters of, 81
 Aix-la-Chapelle waters, 135
 Alkalies, 135, 137
 Alkaline diuretics, 135
 sulphites, 168
 Alopecia, 21
 Alopecia areata. (*See* Tinea
 decalvans)
 Alopecia areata, terminating in
 vitiligo, 17
 Ammonia, carbonate of, 136
 Analysis of 11,000 consecutive
 cases, 3
 Animal parasitic affections, 6, 8,
 32-34
 Aperients, 132
 Arsenic, 143
 Arsenical caustics, 114
 Asiatic pills, 149
 Atrophy of skin, 5, 8, 20

B.

BAREGES, 135
 Baths, 82

Bicarbonate of potash, 136
 Bismuth ointment, 87
 Black soap, 95
 Blisters, 120
 Bloody sweat, 13
 Bromide of potassium, 138

C.

CALOMEL, 98, 134
 Camphor, 85
 Cancer, primary, of skin, 9, 61
 Canities, 24
 Cantharides, 120
 Carbolic acid, 94, 152
 Carbonate of magnesia, 85
 zinc, 85
 ammonia, 136
 Cases of canities, 24
 eczema, 105, 108, 109, 121,
 140, 145
 elephantiasis Græcorum, .59
 ephidrosis cruenta, 14
 furunculi, 172
 hereditary syphilis, 35, 36,
 38
 hyperidrosis, 13
 lichen ruber, 53
 measles displacing eczema,
 76
 multiple fatty tumours, 62
 pemphigus, 147
 primary cancer of skin, 61
 pruritus, 10

- Cases of pruritus senilis, 112
 psoriasis, 106, 141, 150, 154,
 156, 158
 purpura hæmorrhagica, 68
 ,, tuberculosa, 70
 rheumatic purpura, 64, 66
 scleroderma, 56, 138
 strumous disease of toes,
 &c., 111
 syphilitic affection of soles,
 160
 tar acne, 91
 vitiligo, 17, 19
 Causes, Diseases defined by
 uniform, 6, 8, 26-42
 Causes, Diseases not defined by
 uniform, 7, 9, 43
 Causes of eczema, 44
 predisposing, of vegetable
 parasitic affections, 31
 rosacea, 52
 Caustics, 114
 Cautery, actual, 118
 Chronic inflammation of skin,
 90
 Circumscribed eruptions, treat-
 ment of, 120
 Citrate of potash, 136
 Classification of diseases of skin, 4
 10,000 consecutive hospital
 cases, 5
 1,000 consecutive private
 cases, 8
 Clothing, disinfection of, 129
 Cod-liver oil, 138
 Comedones, 5, 8, 12
 Condylomata, treatment of, 98
 Constitutional treatment, 130
 Creasote, 93
 Crusts, removal of, 86
 Disinfection of clothing, 129
 Diuretics, 135
 "Driving in" an eruption,
 danger of, 75
 E.
 ECTHYMA, 7, 47
 Eczema, 7, 9, 43
 displaced by measles, 76
 hereditary syphilitic, 36
 marginatum, 29
 mercuriale, 164
 pruriginosum, 46
 treatment of, 94, 96, 99,
 102, 115, 135
 Elephantiasis Arabum, 7, 9, 58
 Græcorum, 7, 9, 59
 Emplastrum hydrargyri in treat-
 ment of lupus, 99
 Empyreumatic oils, 90, 149
 Enghien, 135
 Ephelis, 5, 8, 16
 Ephidrosis cruenta, 13, 64
 Epilation, 127
 Epithelioma, treatment of, 114
 Erysipelas, treatment of, 116
 Erythema, 7, 9, 45
 treatment of, 94, 98, 135
 F.
 FATTY tumours, 9, 62
 Fevers. (*See* *Tinea favosa*)
 Fevers, eruptive, 6, 41
 Fowler's solution, 148
 Fragilitas crinium, 23
 Functional affections, 5, 8, 9-25
 Furunculi, 172
 G.

D.

DISEASES defined by uniform
 causes, 6, 8, 26-42

GOUTY diathesis, treatment of,
 136

Green iodide of mercury, 98
Guyot's solution of tar, 92

H.

HÆMORRHAGES, 7, 9, 63
Hair, functional diseases of, 5,
8, 21-25
Harrogate waters, 135
Hereditary syphilis, 6, 8, 35-
40
Hereditary character of psoriasis,
48
Hirsuties, 22
Hydropathic treatment, 83
Hyperidrosis, 5, 8, 13
Hyposulphites, 168

I.

IMPERMEABLE dressings, 105°
India-rubber dressings, 105
Infantile syphilis, treatment of,
163
Inflammations, 7, 9, 43-54.
characters of acute, 81
 " chronic, 90
Instruments for actual cautery,
118
epilation, 127
Iodide of potassium, 166
Iodine, 122, 142
Itch. (*See* Scabies)

L.

LEAD ointment, 88
lotion, 88
Lepa, 47
Lice. (*See* Phtheiriasis)
Lichen planus, 53
ruber, 9, 53
scrofulosorum, 6, 8, 41
Liquor carbonis detergens, 92
potassæ, 95

Local treatment, 78
applications act on the
system, 79
applications not uniform in
their action, 80
Lotions of carbonate of soda, 88
of lead, 88
soothing, 88
Luchon, 135
Lupus erythematodes, 8, 41
treatment of, 99, 121
vulgaris, 6, 8, 40
 " treatment of, 104,
116, 118, 121, 122
Lycopodium, 85

M.

MAGNESIA, carbonate of, 85
Melanopathia, 5, 16
Mercurial applications, 98, 118
belt, 164
vapour bath, 162
Mercury, 159
subcutaneous injection of,
161
Microsporon furfur, 29
Miliun, 5, 12
Mineral waters, 135
Moffat waters, 135

N.

NARCOTICS, 137, 138
Neuralgia following shingles,
treatment of, 138
New formations, 7, 9, 55-63
Nitrate of mercury ointment, 98
silver ointment, 116

O.

OIL of Cade, 92
Ointments of bismuth, 87
lead, 88

N

Ointments, mercurial, 98
 of oxide of zinc, 87
 soothing, 86
 Oleum Rusci, 92
 Opium, 138
 Organic affections, 6-9, 26

P.

PARASITIC affections, 6, 8, 26-
 34, 124
 Parasiticides, 132
 Pediculus capitis, 33
 corporis, 33
 pubis, 33
 Pemphigus, 7, 9, 53
 foliaceus, 7, 53
 hereditary syphilitic, 35, 38
 Perchloride of mercury, 96, 98,
 120, 125
 Pfeuffer's treatment, 97
 Phtheiriasis, 6, 8, 33, 46, 125
 Pigmentary disorders of skin, 5,
 8, 16-20
 Pityriasis, 45
 rubra, 7, 9, 45, 53
 Pix liquida, 92, 149
 mineralis, 92, 149
 Potash applications, 95
 effect of, in eczema, 97
 Potash salts, 136
 Potassa fusa, 95, 115
 Potassium, iodide of, 166
 Poultices, 81, 86
 Pruriginous eruption, 46
 Prurigo, 7, 9, 46
 treatment of, 103
 Psoriasis, 7, 9, 47, 154, 156, 158
 in subject of hereditary
 syphilis, 48
 treatment of, 93, 97, 99,
 153
 Purgatives, 132
 Purpura, 7, 9, 64
 pemphigoides, 64
 rheumatica, 64

R.

RED oxide of mercury, 98
 iodide of mercury, 98
 Rheumatic diathesis, treatment
 of, 136
 purpura, 64
 Ringworm. (*See* Tinea trico-
 phytina)
 Rosacea, 7, 9, 51
 treatment of, 52, 102
 Rules for administration of
 arsenic, 144
 iodide of potassium, 166
 mercury, 160

S.

SALIVATION, danger of, from
 external use of mercury, 100
 Sapo mollis, 95
 Scabies, 6, 8, 32, 46
 treatment of, 102, 125
 Scleroderma, 7, 9, 56
 Scrofuloderma, 6, 8, 41
 Sea-water and air, effects of, 85,
 132
 Sebaceous glands, diseases of, 5,
 11-13
 Seborrhœa, 5, 11, 45
 Sedatives, 137
 Shower bath, 84
 Smith's emplastrum canthari-
 dinis liquidum, 120
 Soda, acetate of, 104
 hyposulphite of, 127
 Solution of chlorate of lime, 125
 Soothing applications, 82, 86
 Spring water, danger of, 85
 Staphisagria, ointment of, 125
 Starch powder, 85
 Strumous affections, 6, 8, 40
 treatment of, 104, 122, 138
 diathesis, relation of warts
 to, 56

Styrax ointment, 125
 Subcutaneous injection of mercury, 161
 Sudoriparous glands, diseases of, 5, 8, 13
 Sulphites, 168
 Sulphur, 101, 134
 Sulphurous acid, 127
 Syphilis, hereditary, and psoriasis combined, 49
 Syphilitic affections, 5, 8, 34-40, 160
 condylomata, treatment of, 98
 eruptions, treatment of, 99, 159

T.

TALC, 85
 Tar acne, 91
 Tar rubbed into skin, effect of, 79
 Tar water and pills, 149
 Tarry applications, 92
 Tinea circinata, 6, 8, 29
 decalvans, 6, 8, 30
 ,, treatment of, 122
 favosa, 6, 8, 27
 sycosis, 6, 8, 29
 ,, treatment of, 82
 tonsurans, 6, 8, 29
 ,, treatment of, 121, 123
 tricophytina, 6, 8, 28
 versicolor, 6, 8, 29
 ,, treatment of, 95
 Tonic aperients, 133
 Tonics, 143
 Treatment of acne vulgaris, 50-98, 103
 circumscribed eruptions, 120
 condylomata, 98

Treatment of diseases of the skin, 75
 diseases of the skin, local, 78
 diseases of the skin, constitutional, 130
 ecthyma, 47
 eczema, 94, 96, 99, 102, 108, 109, 115, 121, 132, 135
 epithelioma, 114
 erysipelas, 116
 erythema, 94, 98, 135
 furunculi, 172
 gouty cases, 136
 hirsuties, 22
 ichthyosis, 112
 infantile syphilis, 163
 lupus, 104, 116, 118, 121, 122
 lupus erythematodes, 99, 121, 123
 phtheiriasis, 33, 125
 prurigo, 103, 112
 pruritus senilis, 112
 psoriasis, 49, 93, 95, 99, 106, 150, 153
 rheumatic cases, 136
 rosacea, 52, 102, 150
 scabies, 102, 125
 strumous affections, 41, 104, 111, 122, 138
 syphilitic affections, 34, 99, 159
 tinea decalvans, 122
 ,, sycosis, 82
 ,, tonsurans, 121, 123
 ,, versicolor, 95
 ulcers, 120, 138
 urticaria, 138
 vegetable parasitic affections, 31, 95, 124, 127
 verruca, 56
 zona, 138
 Tricophyton, 28

U.

ULCERS, treatment of, 120
Urticaria, 7, 9, 46

V.

VARIX, 55
Vegetable parasitic affections,
6, 8, 26-32
Verruca, 55
Vitiligo, 15, 17
Vulcanized india-rubber, treat-
ment by means of, 105

W.

WARTS, 55
White precipitate, 98

Z.

ZINC, carbonate of, 85
oxide of, 85
oxide, ointment of, 87
Zona, treatment of, 138

THE END.

BEDFORD STREET, COVENT GARDEN, LONDON,
April 1872.

MACMILLAN & CO.'S CATALOGUE of WORKS
in MATHEMATICS and PHYSICAL SCIENCE;
including PURE and APPLIED MATHE-
MATICS; PHYSICS, ASTRONOMY, GEOLOGY,
CHEMISTRY, ZOOLOGY, BOTANY; PHYSIOLOGY,
ANATOMY and MEDICAL WORKS generally;
and of WORKS in MENTAL and MORAL
PHILOSOPHY and Allied Subjects.

MATHEMATICS.

Airy.—Works by G. B. AIRY, Astronomer Royal :—

ELEMENTARY TREATISE ON PARTIAL DIFFERENTIAL
EQUATIONS. Designed for the Use of Students in the Univer-
sities. With Diagrams. Crown 8vo. cloth. 5s. 6d.

*It is hoped that the methods of solution here explained, and the in-
stances exhibited, will be found sufficient for application to nearly
all the important problems of Physical Science, which require for
their complete investigation the aid of Partial Differential Equa-
tions.*

ON THE ALGEBRAICAL AND NUMERICAL THEORY OF
ERRORS OF OBSERVATIONS AND THE COMBINA-
TION OF OBSERVATIONS. Crown 8vo. cloth. 6s. 6d.

*In order to spare astronomers and observers in natural philosophy the
confusion and loss of time which are produced by referring to the
ordinary treatises embracing both branches of probabilities (the first*

Airy (G. B.)—continual.

relating to chances which can be altered only by the changes of entire units or integral multiples of units in the fundamental conditions of the problem; the other concerning those chances which have respect to insensible gradations in the value of the element measured), this volume has been drawn up. It relates only to errors of observation, and to the rules, derivable from the consideration of these errors, for the combination of the results of observations.

UNDULATORY THEORY OF OPTICS. Designed for the Use of Students in the University. New Edition. Crown 8vo. cloth. 6s. 6d.

The undulatory theory of optics is presented to the reader as having the same claims to his attention as the theory of gravitation,—namely, that it is certainly true, and that, by mathematical operations of general elegance, it leads to results of great interest. This theory explains with accuracy a vast variety of phenomena of the most complicated kind. The plan of this tract has been to include those phenomena only which admit of calculation, and the investigations are applied only to phenomena which actually have been observed.

ON SOUND AND ATMOSPHERIC VIBRATIONS. With the Mathematical Elements of Music. Designed for the Use of Students of the University. Second Edition, revised and enlarged. Crown 8vo. 9s.

This volume consists of sections, which again are divided into numbered articles, on the following topics:—General recognition of the air as the medium which conveys sound; Properties of the air on which the formation and transmission of sound depend; Theory of undulations as applied to sound, etc.; Investigation of the motion of a wave of air through the atmosphere; Transmission of waves of soniferous vibrations through different gases, solids, and fluids; Experiments on the velocity of sound, etc.; On musical sounds, and the manner of producing them; On the elements of musical harmony and melody, and of simple musical composition; On instrumental music; On the human organs of speech and hearing.

A TREATISE ON MAGNETISM. Designed for the Use of Students in the University. Crown 8vo. 9s. 6d.

As the laws of Magnetic Force have been experimentally examined,

with philosophical accuracy, only in its connection with iron and steel, and in the influence excited by the earth as a whole, the accurate portions of this work are confined to the investigations connected with these metals and the earth. The latter part of the work, however, treats in a more general way of the laws of the connection between Magnetism on the one hand and Galvanism and Thermo-Electricity on the other. The work is divided into Twelve Sections, and each section into numbered articles, each of which states concisely and clearly the subject of the following paragraphs.

Ball (R. S., A.M.)—EXPERIMENTAL MECHANICS. A Course of Lectures delivered at the Royal College of Science for Ireland. By ROBERT STAWELL BALL, A.M., Professor of Applied Mathematics and Mechanics in the Royal College of Science for Ireland (Science and Art Department). Royal 8vo. 16s.

The author's aim in these twenty Lectures has been to create in the mind of the student physical ideas corresponding to theoretical laws, and thus to produce a work which may be regarded either as a supplement or an introduction to manuals of theoretic mechanics. To realize this design, the copious use of experimental illustrations was necessary. The apparatus used in the Lectures and figured in the volume has been principally built up from Professor Willis's most admirable system. In the selection of the subjects, the question of practical utility has in many cases been regarded as the one of paramount importance, and it is believed that the mode of treatment which is adopted is more or less original. This is especially the case in the Lectures relating to friction, to the mechanical powers, to the strength of timber and structures, to the laws of motion, and to the pendulum. The illustrations, drawn from the apparatus, are nearly all original and are beautifully executed. "In our reading we have not met with any book of the sort in English."—Mechanics' Magazine.

Bayma.—THE ELEMENTS OF MOLECULAR MECHANICS. By JOSEPH BAYMA, S.J., Professor of Philosophy, Stonyhurst College. Demy 8vo. cloth. 10s. 6d.

Of the twelve Books into which this treatise is divided, the first and second give the demonstration of the principles which bear directly on the constitution and the properties of matter. The next

three books contain a series of theorems and of problems on the laws of motion of elementary substances. In the sixth and seventh, the mechanical constitution of molecules is investigated and determined: and by it the general properties of bodies are explained. The eighth book treats of luminiferous ether. The ninth explains some special properties of bodies. The tenth and eleventh contain a radical and lengthy investigation of chemical principles and relations, which may lead to practical results of high importance. The twelfth and last book treats of molecular masses, distances, and powers.

Boole.—Works by G. BOOLE, D.C.L., F.R.S., Professor of Mathematics in the Queen's University, Ireland :—

A TREATISE ON DIFFERENTIAL EQUATIONS. New and revised Edition. Edited by I. TODHUNTER. Crown 8vo. cloth. 14s.

Professor Boole has endeavoured in this treatise to convey as complete an account of the present state of knowledge on the subject of Differential Equations, as was consistent with the idea of a work intended, primarily, for elementary instruction. The earlier sections of each chapter contain that kind of matter which has usually been thought suitable for the beginner, while the latter ones are devoted either to an account of recent discovery, or the discussion of such deeper questions of principle as are likely to present themselves to the reflective student in connection with the methods and processes of his previous course.

A TREATISE ON DIFFERENTIAL EQUATIONS. Supplementary Volume. Edited by I. TODHUNTER. Crown 8vo. cloth. 8s. 6d.

This volume contains all that Professor Boole wrote for the purpose of enlarging his treatise on Differential Equations.

THE CALCULUS OF FINITE DIFFERENCES. Crown 8vo. cloth. 10s. 6d.

In this exposition of the Calculus of Finite Differences, particular attention has been paid to the connection of its methods with those of the Differential Calculus—a connection which in some instances involves far more than a merely formal analogy. The work is in some measure designed as a sequel to Professor Boole's Treatise on Differential Equations.

Cambridge Senate-House Problems and Riders,
WITH SOLUTIONS:—

1848-1851.—PROBLEMS. By FERRERS and JACKSON. 8vo.
cloth. 15s. 6d.

1848-1851.—RIDERS. By JAMESON. 8vo. cloth. 7s. 6d.

1854.—PROBLEMS AND RIDERS. By WALTON and
MACKENZIE. 8vo. cloth. 10s. 6d.

1857.—PROBLEMS AND RIDERS. By CAMPION and
WALTON. 8vo. cloth. 8s. 6d.

1860.—PROBLEMS AND RIDERS. By WATSON and ROUTH.
Crown 8vo. cloth. 7s. 6d.

1864.—PROBLEMS AND RIDERS. By WALTON and WIL-
KINSON. 8vo. cloth. 10s. 6d.

*These volumes will be found of great value to Teachers and Students,
as indicating the style and range of mathematical study in the
University of Cambridge.*

Cambridge and Dublin Mathematical Journal.

The Complete Work, in Nine Vols. 8vo. cloth. 10l. 10s.

*Only a few copies remain on hand. Among contributors to this
work will be found Sir W. Thomson, Stokes, Adams, Boole, Sir
W. R. Hamilton, De Morgan, Cayley, Sylvester, Jellett, and other
distinguished mathematicians.*

Cheyne.—Works by C. H. H. CHEYNE, M.A., F.R.A.S.:—

AN ELEMENTARY TREATISE ON THE PLANETARY
THEORY. With a Collection of Problems. Second Edition.
Crown 8vo. cloth. 6s. 6d.

*In this volume, an attempt has been made to produce a treatise on the
Planetary theory, which, being elementary in character, should be
so far complete as to contain all that is usually required by students
in the University of Cambridge. This Edition has been carefully
revised. The stability of the Planetary System has been more fully
treated, and an elegant geometrical explanation of the formulæ for
the secular variation of the node and inclination has been in-
troduced.*

THE EARTH'S MOTION OF ROTATION. Crown 8vo.
3s. 6d.

The first part of this work consists of an application of the method of the variation of elements to the general problem of rotation. In the second part the general rotation formulæ are applied to the particular case of the earth.

Childe.—THE SINGULAR PROPERTIES OF THE ELLIPSOID AND ASSOCIATED SURFACES OF THE N^{TH} DEGREE. By the Rev. G. F. CHILDE, M.A., Author of "Ray Surfaces," "Related Caustics," &c. 8vo. 10s. 6d.

The object of this volume is to develop peculiarities in the Ellipsoid; and, further, to establish analogous properties in the unlimited congeneric series of which this remarkable surface is a constituent.

Dodgson.—AN ELEMENTARY TREATISE ON DETERMINANTS, with their Application to Simultaneous Linear Equations and Algebraical Geometry. By CHARLES L. DODGSON, M.A., Student and Mathematical Lecturer of Christ Church, Oxford. Small 4to. cloth. 10s. 6d.

The object of the author is to present the subject as a continuous chain of argument, separated from all accessories of explanation or illustration. All such explanation and illustration as seemed necessary for a beginner are introduced either in the form of foot-notes, or, where that would have occupied too much room, of Appendices.

Earnshaw (S., M.A.)—PARTIAL DIFFERENTIAL EQUATIONS. An Essay towards an entirely New Method of Integrating them. By S. EARNSHAW, M.A., of St. John's College, Cambridge. Crown 8vo. 5s.

The peculiarity of the system expounded in this work is, that in every equation, whatever be the number of original independent variables, the work of integration is at once reduced to the use of one independent variable only. The author's object is merely to render his method thoroughly intelligible. The various steps of the investigation are all obedient to one general principle: and though in some degree novel, are not really difficult, but on the contrary easy when the eye has become accustomed to the novelties of the notation. Many of the results of the integrations are far more general than they were in the shape in which they appeared in former Treatises, and many

Equations will be found in this Essay integrated with ease in finite terms, which were never so integrated before.

Ferrers.—AN ELEMENTARY TREATISE ON TRILINEAR CO-ORDINATES, the Method of Reciprocal Polars, and the Theory of Projectors. By the Rev. N. M. FERRERS, M.A., Fellow and Tutor of Gonville and Caius College, Cambridge. Second Edition. Crown 8vo. 6s. 6d.

The object of the author in writing on this subject has mainly been to place it on a basis altogether independent of the ordinary Cartesian system, instead of regarding it as only a special form of Abridged Notation. A short chapter on Determinants has been introduced.

Frost.—THE FIRST THREE SECTIONS OF NEWTON'S PRINCIPIA. With Notes and Illustrations. Also a Collection of Problems, principally intended as Examples of Newton's Methods. By PERCIVAL FROST, M.A., late Fellow of St. John's College, Mathematical Lecturer of King's College, Cambridge. Second Edition. 8vo. cloth. 10s. 6d.

The author's principal intention is to explain difficulties which may be encountered by the student on first reading the Principia, and to illustrate the advantages of a careful study of the methods employed by Newton, by showing the extent to which they may be applied in the solution of problems; he has also endeavoured to give assistance to the student who is engaged in the study of the higher branches of mathematics, by representing in a geometrical form several of the processes employed in the Differential and Integral Calculus, and in the analytical investigations of Dynamics.

Frost and Wolstenholme.—A TREATISE ON SOLID GEOMETRY. By PERCIVAL FROST, M.A., and the Rev. J. WOLSTENHOLME, M.A., Fellow and Assistant Tutor of Christ's College. 8vo. cloth. 18s.

Intending to make the subject accessible, at least in the earlier portions to all classes of students, the authors have endeavoured to explain completely all the processes which are most useful in dealing with ordinary theorems and problems, thus directing the student to the selection of methods which are best adapted to the exigencies of each problem. In the more difficult portions of the subject, they have

considered themselves to be addressing a higher class of students ; and they have there tried to lay a good foundation on which to build, if any reader should wish to pursue the science beyond the limits to which the work extends.

Godfray.—Works by HUGH GODFRAY, M.A., Mathematical Lecturer at Pembroke College, Cambridge :—

A TREATISE ON ASTRONOMY, for the Use of Colleges and Schools. 8vo. cloth. 12s. 6d.

This book embraces all those branches of Astronomy which have, from time to time, been recommended by the Cambridge Board of Mathematical Studies : but by far the larger and easier portion, adapted to the first three days of the Examination for Honours, may be read by the more advanced pupils in many of our schools. The author's aim has been to convey clear and distinct ideas of the celestial phenomena. "It is a working book," says the Guardian, "taking Astronomy in its proper place in the Mathematical Sciences. . . . It is a book which is not likely to be got up unintelligently."

AN ELEMENTARY TREATISE ON THE LUNAR THEORY, with a Brief Sketch of the Problem up to the time of Newton. Second Edition, revised. Crown 8vo. cloth. 5s. 6d.

These pages will, it is hoped, form an introduction to more recondite works. Difficulties have been discussed at considerable length. The selection of the method followed with regard to analytical solutions, which is the same as that of Airy, Herschell, etc., was made on account of its simplicity ; it is, moreover, the method which has obtained in the University of Cambridge. "As an elementary treatise and introduction to the subject, we think it may justly claim to supersede all former ones."—London, Edinburgh, and Dublin Phil. Magazine.

Green (George).—MATHEMATICAL PAPERS OF THE LATE GEORGE GREEN, Fellow of Gonville and Caius College, Cambridge. Edited by N. M. FERRERS, M.A., Fellow and Tutor of Gonville and Caius College. 8vo. 15s.

The publication of this book may be opportune at present, as several of the subjects with which they are directly or indirectly concerned have recently been introduced into the course of mathematical

study at Cambridge. They have also an interest as being the work of an almost entirely self-taught mathematical genius. The Papers comprise the following:—An Essay on the application of Mathematical Analysis to the Theories of Electricity and Magnetism—On the Laws of the Equilibrium of Fluids analogous to the Electric Fluid—On the Determination of the Attractions of Ellipsoids of variable Densities—On the Motion of Waves in a variable Canal of small depth and width—On the Reflection and Refraction of Sound—On the Reflection and Refraction of Light at the Common Surface of two Non-Crystallized Media—On the Propagation of Light in Crystallized Media—Researches on the Vibrations of Pendulums in Fluid Media. “It has been for some time recognized that Green’s writings are amongst the most valuable mathematical productions we possess.”—Athenæum.

Hemming.—AN ELEMENTARY TREATISE ON THE DIFFERENTIAL AND INTEGRAL CALCULUS. For the Use of Colleges and Schools. By G. W. HEMMING, M.A., Fellow of St. John’s College, Cambridge. Second Edition, with Corrections and Additions. 8vo. cloth. 9s.

“There is no book in common use from which so clear and exact a knowledge of the principles of the Calculus can be so readily obtained.”—Literary Gazette.

Jackson.—GEOMETRICAL CONIC SECTIONS. An Elementary Treatise in which the Conic Sections are defined as the Plane Sections of a Cone, and treated by the Method of Projections. By J. STUART JACKSON, M.A., late Fellow of Gonville and Caius College. Crown 8vo. 4s. 6d.

This work has been written with a view to give the student the benefit of the Method of Projections as applied to the Ellipse and Hyperbola. When this method is admitted into the treatment of Conic Sections there are many reasons why they should be defined, not with reference to the focus and directrix, but according to the original definition from which they have their name, as Plane Sections of a Cone. This method is calculated to produce a material simplification in the treatment of these curves and to make the proof of their properties more easily understood in the first instance and more easily remembered. It is also a powerful instrument in the solution of a large class of problems relating to these curves.

Morgan.—A COLLECTION OF PROBLEMS AND EXAMPLES IN MATHEMATICS. With Answers. By H. A. MORGAN, M.A., Sadlerian and Mathematical Lecturer of Jesus College, Cambridge. Crown 8vo. cloth. 6s. 6d.

This book contains a number of problems, chiefly elementary, in the Mathematical subjects usually read at Cambridge. They have been selected from the Papers set during late years at Jesus College. Very few of them are to be met with in other collections, and by far the larger number are due to some of the most distinguished Mathematicians in the University.

Newton's Principia.—4to. cloth. 31s. 6d.

It is a sufficient guarantee of the reliability of this complete edition of Newton's Principia that it has been printed for and under the care of Professor Sir William Thomson and Professor Blackburn, of Glasgow University. The following notice is prefixed:—"Finding that all the editions of the Principia are now out of print, we have been induced to reprint Newton's last edition [of 1726] without note or comment, only introducing the 'Corrigenda' of the old copy and correcting typographical errors." The book is of a handsome size, with large type, fine thick paper, and cleanly-cut figures, and is the only recent edition containing the whole of Newton's great work.

Parkinson.—Works by S. PARKINSON, D.D., F.R.S., Fellow and Tutor of St. John's College, Cambridge :—

AN ELEMENTARY TREATISE ON MECHANICS. For the Use of the Junior Classes at the University and the Higher Classes in Schools. With a Collection of Examples. Fourth Edition, revised. Crown 8vo. cloth. 9s. 6d.

In preparing a fourth edition of this work the author has kept the same object in view as he had in the former editions—namely, to include in it such portions of Theoretical Mechanics as can be conveniently investigated without the use of the Differential Calculus, and so render it suitable as a manual for the junior classes in the University and the higher classes in Schools. With one or two short exceptions, the student is not presumed to require a knowledge of any branches of Mathematics beyond the elements of Algebra, Geometry,

Parkinson (S.)—*continued.*

and Trigonometry. Several additional propositions have been incorporated in the work for the purpose of rendering it more complete, and the collection of Examples and Problems has been largely increased.

A TREATISE ON OPTICS. Third Edition, revised and enlarged. Crown 8vo. cloth. 10s. 6d.

A collection of Examples and Problems has been appended to this work, which are sufficiently numerous and varied in character to afford useful exercise for the student. For the greater part of them, recourse has been had to the Examination Papers set in the University and the several Colleges during the last twenty years.

Phear.—ELEMENTARY HYDROSTATICS. With Numerous Examples. By J. B. PHEAR, M.A., Fellow and late Assistant Tutor of Clare College, Cambridge. Fourth Edition. Crown 8vo. cloth. 5s. 6d.

This edition has been carefully revised throughout, and many new Illustrations and Examples added, which it is hoped will increase its usefulness to students at the Universities and in Schools. In accordance with suggestions from many engaged in tuition, answers to all the Examples have been given at the end of the book.

Pratt.—A TREATISE ON ATTRACTIONS, LAPLACE'S FUNCTIONS, AND THE FIGURE OF THE EARTH. By JOHN H. PRATT, M.A., Archdeacon of Calcutta, Author of "The Mathematical Principles of Mechanical Philosophy." Fourth Edition. Crown 8vo. cloth. 6s. 6d.

The author's chief design in this treatise is to give an answer to the question, "Has the Earth acquired its present form from being originally in a fluid state?" This edition is a complete revision of the former ones.

Puckle.—AN ELEMENTARY TREATISE ON CONIC SECTIONS AND ALGEBRAIC GEOMETRY. With numerous Examples and Hints for their Solution; especially designed for the Use of Beginners. By G. H. PUCKLE, M.A., Head Master of

Windermere College. New Edition, revised and enlarged. Crown 8vo. cloth. 7s. 6d.

This work is recommended by the Syndicate of the Cambridge Local Examinations, and is the text-book in Harvard University, U.S. The Athenæum says the author "displays an intimate acquaintance with the difficulties likely to be felt, together with a singular aptitude in removing them."

Routh.—AN ELEMENTARY TREATISE ON THE DYNAMICS OF THE SYSTEM OF RIGID BODIES. With numerous Examples. By EDWARD JOHN ROUTH, M.A., late Fellow and Assistant Tutor of St. Peter's College, Cambridge; Examiner in the University of London. Second Edition, enlarged. Crown 8vo. cloth. 14s.

In this edition the author has made several additions to each chapter: he has tried, even at the risk of some little repetition, to make each chapter, as far as possible, complete in itself, so that all that relates to any one part of the subject may be found in the same place. This arrangement will enable every student to select his own order in which to read the subject. The Examples which will be found at the end of each chapter have been chiefly selected from the Examination Papers which have been set in the University and the Colleges in the last few years.

Smith's (Barnard) Works.—See EDUCATIONAL CATALOGUE.

Smith (J. Brook.)—ARITHMETIC IN THEORY AND PRACTICE. By J. BROOK SMITH, M.A., LL.B., St. John's College, Cambridge; Barrister-at-Law; one of the Masters of Cheltenham College. Crown 8vo. 4s. 6d.

Writers on Arithmetic at the present day feel the necessity of explaining the principles on which the rules of the subject are based, but few as yet feel the necessity of making these explanations strict and complete; or, failing that, of distinctly pointing out their defective character. If the science of Arithmetic is to be made an effective instrument in developing and strengthening the mental powers, it ought to be worked out rationally and conclusively; and in this work the author has endeavoured to reason out in a clear and accurate

manner the leading propositions of the science, and to illustrate and apply those propositions in practice. In the practical part of the subject he has advanced somewhat beyond the majority of preceding writers; particularly in Division, in Greatest Common Measure, in Cube Root, in the chapters on Decimal Money and the Metric System, and more especially in the application of Decimals to Percentages and cognate subjects. Copious examples, original and selected, are given.

Snowball.—THE ELEMENTS OF PLANE AND SPHERICAL TRIGONOMETRY; with the Construction and Use of Tables of Logarithms. By J. C. SNOWBALL, M.A. Tenth Edition. Crown 8vo. cloth. 7s. 6d.

In preparing the present edition for the press, the text has been subjected to a careful revision; the proofs of some of the more important propositions have been rendered more strict and general; and a considerable addition of more than two hundred examples, taken principally from the questions set of late years in the public examinations of the University and of individual Colleges, has been made to the collection of Examples and Problems for practice.

Tait and Steele.—DYNAMICS OF A PARTICLE. With numerous Examples. By Professor TAIT and Mr. STEELE. New Edition. Crown 8vo. cloth. 10s. 6d.

In this treatise will be found all the ordinary propositions, connected with the Dynamics of Particles, which can be conveniently deduced without the use of D'Alembert's Principle. Throughout the book will be found a number of illustrative examples introduced in the text, and for the most part completely worked out; others with occasional solutions or hints to assist the student are appended to each chapter. For by far the greater portion of these, the Cambridge Senate-House and College Examination Papers have been applied to.

Taylor.—GEOMETRICAL CONICS; including Anharmonic Ratio and Projection, with numerous Examples. By C. TAYLOR, B.A., Scholar of St. John's College, Cambridge. Crown 8vo. cloth. 7s. 6d.

This work contains elementary proofs of the principal properties of Conic Sections, together with chapters on Projection and Anharmonic Ratio.

Todhunter.—Works by I. TODHUNTER, M.A., F.R.S., of St. John's College, Cambridge :—

"Perspicuous language, vigorous investigations, scrutiny of difficulties, and methodical treatment, characterize Mr. Todhunter's works."—Civil Engineer.

THE ELEMENTS OF EUCLID; MENSURATION FOR BEGINNERS; ALGEBRA FOR BEGINNERS; TRIGONOMETRY FOR BEGINNERS; MECHANICS FOR BEGINNERS.—See EDUCATIONAL CATALOGUE.

ALGEBRA. For the Use of Colleges and Schools. Fifth Edition. Crown 8vo. cloth. 7s. 6d.

This work contains all the propositions which are usually included in elementary treatises on Algebra, and a large number of Examples for Exercise. The author has sought to render the work easily intelligible to students, without impairing the accuracy of the demonstrations, or contracting the limits of the subject. The Examples, about Sixteen hundred and fifty in number, have been selected with a view to illustrate every part of the subject. The work will be found peculiarly adapted to the wants of students who are without the aid of a teacher. The Answers to the Examples, with hints for the solution of some in which assistance may be needed, are given at the end of the book. In the present edition two New Chapters and Three hundred miscellaneous Examples have been added. "It has merits which unquestionably place it first in the class to which it belongs."—Educator.

KEY TO ALGEBRA FOR THE USE OF COLLEGES AND SCHOOLS. Crown 8vo. 10s. 6d.

AN ELEMENTARY TREATISE ON THE THEORY OF EQUATIONS. Second Edition, revised. Crown 8vo. cloth. 7s. 6d.

This treatise contains all the propositions which are usually included in elementary treatises on the theory of Equations, together with Examples for exercise. These have been selected from the College and University Examination Papers, and the results have been given when it appeared necessary. In order to exhibit a comprehensive view of the subject, the treatise includes investigations which are not found in all the preceding elementary treatises, and also

Todhunter (I.)—continued.

some investigations which are not to be found in any of them. For the second edition the work has been revised and some additions have been made, the most important being an account of the Researches of Professor Sylvester respecting Newton's Rule. "A thoroughly trustworthy, complete, and yet not too elaborate treatise."
—Philosophical Magazine.

PLANE TRIGONOMETRY. For Schools and Colleges. Fourth Edition. Crown 8vo. cloth. 5s.

The design of this work has been to render the subject intelligible to beginners, and at the same time to afford the student the opportunity of obtaining all the information which he will require on this branch of Mathematics. Each chapter is followed by a set of Examples: those which are entitled Miscellaneous Examples, together with a few in some of the other sets, may be advantageously reserved by the student for exercise after he has made some progress in the subject. In the Second Edition the hints for the solution of the Examples have been considerably increased.

A TREATISE ON SPHERICAL TRIGONOMETRY. Third Edition, enlarged. Crown 8vo. cloth. 4s. 6d.

The present work is constructed on the same plan as the treatise on Plane Trigonometry, to which it is intended as a sequel. In the account of Napier's Rules of circular parts, an explanation has been given of a method of proof devised by Napier, which seems to have been overlooked by most modern writers on the subject. Considerable labour has been bestowed on the text in order to render it comprehensive and accurate, and the Examples (selected chiefly from College Examination Papers) have all been carefully verified. "For educational purposes this work seems to be superior to any others on the subject."—Critic.

PLANE CO-ORDINATE GEOMETRY, as applied to the Straight Line and the Conic Sections. With numerous Examples. Fourth Edition, revised and enlarged. Crown 8vo. cloth. 7s. 6d.

The author has here endeavoured to exhibit the subject in a simple manner for the benefit of beginners, and at the same time to include in one volume all that students usually require. In addition, therefore, to the propositions which have always appeared in such

Todhunter (I.)—continued.

treatises, he has introduced the methods of abridged notation, which are of more recent origin: these methods, which are of a less elementary character than the rest of the work, are placed in separate chapters, and may be omitted by the student at first.

A TREATISE ON THE DIFFERENTIAL CALCULUS.

With numerous Examples. Fifth Edition. Crown 8vo. cloth. 10s. 6d.

The author has endeavoured in the present work to exhibit a comprehensive view of the Differential Calculus on the method of limits. In the more elementary portions he has entered into considerable detail in the explanations, with the hope that a reader who is without the assistance of a tutor may be enabled to acquire a competent acquaintance with the subject. The method adopted is that of Differential Coefficients. To the different chapters are appended Examples sufficiently numerous to render another book unnecessary; these Examples being mostly selected from College Examination Papers. This and the following work have been translated into Italian by Professor Battaglini, who in his Preface speaks thus:—
"In publishing this translation of the Differential and Integral Calculus of Mr. Todhunter, we have had no other object than to add to the books which are in the hands of the students of our Universities, a work remarkable for the clearness of the exposition, the rigour of the demonstrations, the just proportion in the parts, and the rich store of examples which offer a large field for useful exercise."

A TREATISE ON THE INTEGRAL CALCULUS AND ITS APPLICATIONS.

With numerous Examples. Third Edition, revised and enlarged. Crown 8vo. cloth. 10s. 6d.

This is designed as a work at once elementary and complete, adapted for the use of beginners, and sufficient for the wants of advanced students. In the selection of the propositions, and in the mode of establishing them, it has been sought to exhibit the principles clearly, and to illustrate all their most important results. The process of summation has been repeatedly brought forward, with the view of securing the attention of the student to the notions which form the true foundation of the Calculus itself, as well as of its most valuable applications. Every attempt has been made to explain those

Todhunter (I.)—*continued.*

difficulties which usually perplex beginners, especially with reference to the limits of integrations. A new method has been adopted in regard to the transformation of multiple integrals. The last chapter deals with the Calculus of Variations. A large collection of Exercises, selected from College Examination Papers, has been appended to the several chapters.

EXAMPLES OF ANALYTICAL GEOMETRY OF THREE DIMENSIONS. Second Edition, revised. Crown 8vo. cloth. 4s.

A TREATISE ON ANALYTICAL STATICS. With numerous Examples. Third Edition, revised and enlarged. Crown 8vo. cloth. 10s. 6d.

In this work on Statics (treating of the laws of the equilibrium of bodies) will be found all the propositions which usually appear in treatises on Theoretical Statics. To the different chapters Examples are appended, which have been principally selected from University Examination Papers. In the Third Edition many additions have been made, in order to illustrate the application of the principles of the subject to the solution of problems.

A HISTORY OF THE MATHEMATICAL THEORY OF PROBABILITY, from the Time of Pascal to that of Laplace. 8vo. 18s.

The subject of this work has high claims to consideration on account of the subtle problems which it involves, the valuable contributions to analysis which it has produced, its important practical applications, and the eminence of those who have cultivated it; nearly every great mathematician within the range of a century and a half comes under consideration in the course of the history. The author has endeavoured to be quite accurate in his statements, and to reproduce the essential elements of the original works which he has analysed. Besides being a history, the work may claim the title of a comprehensive treatise on the Theory of Probability, for it assumes in the reader only so much knowledge as can be gained from an elementary book on Algebra, and introduces him to almost every process and every special problem which the literature of the subject can furnish.

RESEARCHES IN THE CALCULUS OF VARIATIONS,
Principally on the Theory of Discontinuous Solutions: An Essay

Todhunter (I.)—*continued.*

to which the Adams' Prize was awarded in the University of Cambridge in 1871. 8vo. 6s.

The subject of this Essay was prescribed in the following terms by the Examiners :—"A determination of the circumstances under which discontinuity of any kind presents itself in the solution of a problem of maximum or minimum in the Calculus of Variations, and applications to particular instances. It is expected that the discussion of the instances should be exemplified as far as possible geometrically, and that attention be especially directed to cases of real or supposed failure of the Calculus." While the Essay is thus mainly devoted to the consideration of discontinuous solutions, various other questions in the Calculus of Variations are examined and elucidated; and the author hopes he has definitely contributed to the extension and improvement of our knowledge of this refined department of analysis.

Wilson (W. P.)—A TREATISE ON DYNAMICS. By W. P. WILSON, M.A., Fellow of St. John's College, Cambridge, and Professor of Mathematics in Queen's College, Belfast. 8vo. 9s. 6d.

Wolstenholme.—A BOOK OF MATHEMATICAL PROBLEMS, on Subjects included in the Cambridge Course. By JOSEPH WOLSTENHOLME, Fellow of Christ's College, some time Fellow of St. John's College, and lately Lecturer in Mathematics at Christ's College. Crown 8vo. cloth. 8s. 6d.

CONTENTS :—*Geometry (Euclid)—Algebra—Plane Trigonometry—Geometrical Conic Sections—Analytical Conic Sections—Theory of Equations—Differential Calculus—Integral Calculus—Solid Geometry—Statics—Elementary Dynamics—Newton—Dynamics of a Point—Dynamics of a Rigid Body—Hydrostatics—Geometrical Optics—Spherical Trigonometry and Plane Astronomy. In some cases the author has prefixed to certain classes of problems fragmentary notes on the mathematical subjects to which they relate. "Judicious, symmetrical, and well arranged."*—Guardian.

PHYSICAL SCIENCE.

Airy (G. B.)—POPULAR ASTRONOMY. With Illustrations.
By G. B. AIRY, Astronomer Royal. Seventh and cheaper Edition.
18mo. cloth. 4s. 6d.

This work consists of Six Lectures, which are intended "to explain to intelligent persons the principles on which the instruments of an Observatory are constructed (omitting all details, so far as they are merely subsidiary), and the principles on which the observations made with these instruments are treated for deduction of the distances and weights of the bodies of the Solar System, and of a few stars, omitting all minutiae of formulæ, and all troublesome details of calculation." The speciality of this volume is the direct reference of every step to the Observatory, and the full description of the methods and instruments of observation.

Bastian (H. C. M.D., F.R.S.)—THE MODES OF ORIGIN OF LOWEST ORGANISMS: Including a Discussion of the Experiments of M. Pasteur, and a reply to some Statements by Professors Huxley and Tyndall. By H. CHARLTON BASTIAN, M.D., F.R.S., Professor of Pathological Anatomy in University College, London, etc. Crown 8vo. 4s. 6d.

The present volume contains a fragment of the evidence which will be embodied in a much larger work—now almost completed—relating to the nature and origin of living matter, and in favour of what is termed the Physical Doctrine of Life. "It is a work worthy of the highest respect, and places its author in the very first class of scientific physicians. . . . It would be difficult to name an instance in which skill, knowledge, perseverance, and great reasoning power have been more happily applied to the investigation of a complex biological problem."—British Medical Journal.

Birks (R. B.)—ON MATTER AND ETHER ; or, The Secret Laws of Physical Change. By THOMAS RAWSON BIRKS, M.A., Rector of Kelshall, Herts, formerly Fellow of Trinity College, Cambridge. Crown 8vo. 5s. 6d.

The author believes that the hypothesis of the existence of, besides matter, a luminous ether, of immense elastic force, supplies the true and sufficient key to the remaining secrets of inorganic matter, of the phenomena of light, electricity, etc. In this treatise the author endeavours first to form a clear and definite conception with regard to the real nature both of matter and ether, and the laws of mutual action which must be supposed to exist between them. He then endeavours to trace out the main consequences of the fundamental hypothesis, and their correspondence with the known phenomena of physical change.

Blanford (W. T.)—GEOLOGY AND ZOOLOGY OF ABYSSINIA. By W. T. BLANFORD. 8vo. 21s.

This work contains an account of the Geological and Zoological Observations made by the author in Abyssinia, when accompanying the British Army on its march to Magdala and back in 1868, and during a short journey in Northern Abyssinia, after the departure of the troops. Part I. Personal Narrative; Part II. Geology; Part III. Zoology. With Coloured Illustrations and Geological Map. "The result of his labours," the Academy says, "is an important contribution to the natural history of the country."

Cooke (Josiah P., Jun.)—FIRST PRINCIPLES OF CHEMICAL PHILOSOPHY. By JOSIAH P. COOKE, Jun., Ervine Professor of Chemistry and Mineralogy in Harvard College. Crown 8vo. 12s.

The object of the author in this book is to present the philosophy of Chemistry in such a form that it can be made with profit the subject of College recitations, and furnish the teacher with the means of testing the student's faithfulness and ability. With this view the subject has been developed in a logical order, and the principles of the science are taught independently of the experimental evidence on which they rest.

Cooke (M. C.)—HANDBOOK OF BRITISH FUNGI, with full descriptions of all the Species, and Illustrations of the Genera. By M. C. COOKE, M.A. Two vols. crown 8vo. 24s.

During the thirty-five years that have elapsed since the appearance of the last complete Mycologic Flora no attempt has been made to revise it, to incorporate species since discovered, and to bring it up to the standard of modern science. No apology, therefore, is necessary for the present effort, since all will admit that the want of such a manual has long been felt, and this work makes its appearance under the advantage that it seeks to occupy a place which has long been vacant. No effort has been spared to make the work worthy of confidence, and, by the publication of an occasional supplement, it is hoped to maintain it for many years as the "Handbook" for every student of British Fungi. Appended is a complete alphabetical Index of all the divisions and subdivisions of the Fungi noticed in the text. The book contains 400 figures. "Will maintain its place as the standard English book, on the subject of which it treats, for many years to come."—Standard.

Dawson (J. W.)—ACADIAN GEOLOGY. The Geologic Structure, Organic Remains, and Mineral Resources of Nova Scotia, New Brunswick, and Prince Edward Island. By JOHN WILLIAM DAWSON, M.A., LL.D., F.R.S., F.G.S., Principal and Vice-Chancellor of M'Gill College and University, Montreal, &c. Second Edition, revised and enlarged. With a Geological Map and numerous Illustrations. 8vo. 18s.

The object of the first edition of this work was to place within the reach of the people of the districts to which it relates, a popular account of the more recent discoveries in the geology and mineral resources of their country, and at the same time to give to geologists in other countries a connected view of the structure of a very interesting portion of the American Continent, in its relation to general and theoretical Geology. In the present edition, it is hoped this design is still more completely fulfilled, with reference to the present more advanced condition of knowledge. The author has endeavoured to convey a knowledge of the structure and fossils of the region in such a manner as to be intelligible to ordinary readers, and has devoted much attention to all questions relating to the nature and present or prospective value of deposits of useful minerals.

Besides a large coloured Geological Map of the district, the work is illustrated by upwards of 260 cuts of sections, fossils, animals, etc. "The book will doubtless find a place in the library, not only of the scientific geologist, but also of all who are desirous of the industrial progress and commercial prosperity of the Acadian provinces."—Mining Journal. "A style at once popular and scientific. . . . A valuable addition to our store of geological knowledge."—Guardian.

Flower (W. H.)—AN INTRODUCTION TO THE OSTEOLOGY OF THE MAMMALIA. Being the substance of the Course of Lectures delivered at the Royal College of Surgeons of England in 1870. By W. H. FLOWER, F.R.S., F.R.C.S., Hunterian Professor of Comparative Anatomy and Physiology. With numerous Illustrations. Globe 8vo. 7s. 6d.

Although the present work contains the substance of a Course of Lectures, the form has been changed, so as the better to adapt it as a hand-book for students. Theoretical views have been almost entirely excluded: and while it is impossible in a scientific treatise to avoid the employment of technical terms, it has been the author's endeavour to use no more than absolutely necessary, and to exercise due care in selecting only those that seem most appropriate, or which have received the sanction of general adoption. With a very few exceptions the illustrations have been drawn expressly for this work from specimens in the Museum of the Royal College of Surgeons.

Galton.—Works by FRANCIS GALTON, F.R.S. :—

METEOROGRAPHICA, or Methods of Mapping the Weather. Illustrated by upwards of 600 Printed Lithographic Diagrams. 4to. 9s.

As Mr. Galton entertains strong views on the necessity of Meteorological Charts and Maps, he determined, as a practical proof of what could be done, to chart the entire area of Europe, so far as meteorological stations extend, during one month, viz. the month of December, 1861. Mr. Galton got his data from authorities in every part of Britain and the Continent, and on the basis of these has here drawn up nearly a hundred different Maps and Charts, showing the state of the weather all over Europe during the above period. "If the various Governments and scientific bodies would perform for the

Galton (F.)—*continued.*

whole world for two or three years what, at a great cost and labour, Mr. Galton has done for a part of Europe for one month, Meteorology would soon cease to be made a joke of.—Spectator.

HEREDITARY GENIUS: An Inquiry into its Laws and Consequences. Demy 8vo. 12s.

"I propose," the author says, "to show in this book that a man's natural abilities are derived by inheritance, under exactly the same limitations as are the form and physical features of the whole organic world. I shall show that social agencies of an ordinary character, whose influences are little suspected, are at this moment working towards the degradation of human nature, and that others are working towards its improvement. The general plan of my argument is to show that high reputation is a pretty accurate test of high ability; next, to discuss the relationships of a large body of fairly eminent men, and to obtain from these a general survey of the laws of heredity in respect of genius. Then will follow a short chapter, by way of comparison, on the hereditary transmission of physical gifts, as deduced from the relationships of certain classes of oarsmen and wrestlers. Lastly, I shall collate my results and draw conclusions." The Times calls it "a most able and most interesting book;" and Mr. Darwin, in his "Descent of Man" (vol. i. p. 111), says, "We know, through the admirable labours of Mr. Galton, that Genius tends to be inherited."

Geikie (A.)—SCENERY OF SCOTLAND, Viewed in Connection with its Physical Geography. With Illustrations and a new Geological Map. By ARCHIBALD GEIKIE, Professor of Geology in the University of Edinburgh. Crown 8vo. 10s. 6d.

"We can confidently recommend Mr. Geikie's work to those who wish to look below the surface and read the physical history of the Scenery of Scotland by the light of modern science."—Saturday Review.

"Amusing, picturesque, and instructive."—Times.

Hooker (Dr.)—THE STUDENT'S FLORA OF THE BRITISH ISLANDS. By J. D. HOOKER, C.B., F.R.S., M.D., D.C.L., Director of the Royal Gardens, Kew. Globe 8vo. 10s. 6d.

The object of this work is to supply students and field-botanists with a fuller account of the Plants of the British Islands than the manuals hitherto in use aim at giving. The Ordinal, Generic, and Specific characters have been re-written, and are to a great extent original, and drawn from living or dried specimens, or both. "Cannot fail to perfectly fulfil the purpose for which it is intended."—Land and Water. "Containing the fullest and most accurate manual of the kind that has yet appeared."—Pall Mall Gazette.

Huxley (Professor).—LAY SERMONS, ADDRESSES, AND REVIEWS. By T. H. HUXLEY, LL.D., F.R.S. New and Cheaper Edition. Crown 8vo. 7s. 6d.

Fourteen Discourses on the following subjects:—(1) On the Advisableness of Improving Natural Knowledge:—(2) Emancipation—Black and White:—(3) A Liberal Education, and where to find it:—(4) Scientific Education:—(5) On the Educational Value of the Natural History Sciences:—(6) On the Study of Zoology:—(7) On the Physical Basis of Life:—(8) The Scientific Aspects of Positivism:—(9) On a Piece of Chalk:—(10) Geological Contemporaneity and Persistent Types of Life:—(11) Geological Reform:—(12) The Origin of Species:—(13) Criticisms on the "Origin of Species."—(14) On Descartes' "Discourse touching the Method of using One's Reason rightly and of seeking Scientific Truth." The momentous influence exercised by Mr. Huxley's writings on physical, mental, and social science is universally acknowledged: his works must be studied by all who would comprehend the various drifts of modern thought.

ESSAYS SELECTED FROM LAY SERMONS, ADDRESSES, AND REVIEWS. Crown 8vo. 1s.

This volume includes Numbers 1, 3, 4, 7, 8, and 14, of the above.

LESSONS IN ELEMENTARY PHYSIOLOGY. With numerous Illustrations. Fourteenth Thousand. 18mo. cloth. 4s. 6d.

This book describes and explains, in a series of graduated lessons, the principles of Human Physiology, or the Structure and Functions of the Human Body. The first lesson supplies a general view of the subject. This is followed by sections on the Vascular or Venous System, and the Circulation; the Blood and the Lymph; Respiration.

tion : Sources of Loss and of Gain to the Blood ; the Function of Alimentation ; Motion and Locomotion ; Sensations and Sensory Organs ; the Organ of Sight ; the Coalescence of Sensations with one another and with other States of Consciousness ; the Nervous System and Innervation ; Histology, or the Minute Structure of the Tissues. A Table of Anatomical and Physiological Constants is appended. The lessons are fully illustrated by numerous engravings. "Pure gold throughout."—Guardian. "Unquestionably the clearest and most complete elementary treatise on this subject that we possess in any language."—Westminster Review.

Kirchhoff (G.)—RESEARCHES ON THE SOLAR SPECTRUM, and the Spectra of the Chemical Elements. By G. KIRCHHOFF, Professor of Physics in the University of Heidelberg. Second Part. Translated, with the Author's Sanction, from the Transactions of the Berlin Academy for 1862, by HENRY R. ROSCOE, B.A., Ph.D., F.R.S., Professor of Chemistry in Owens College, Manchester.

"It is to Kirchhoff we are indebted for by far the best and most accurate observations of these phenomena."—Edin. Review. "This memoir seems almost indispensable to every Spectrum observer."—Philosophical Magazine.

Lockyer (J. N.)—ELEMENTARY LESSONS IN ASTRONOMY. With numerous Illustrations. By J. NORMAN LOCKYER, F.R.S. Eighth Thousand. 18mo. 5s. 6d.

The author has here aimed to give a connected view of the whole subject, and to supply facts, and ideas founded on the facts, to serve as a basis for subsequent study and discussion. The chapters treat of the Stars and Nebulæ ; the Sun ; the Solar System ; Apparent Movements of the Heavenly Bodies ; the Measurement of Time ; Light ; the Telescope and Spectroscope ; Apparent Places of the Heavenly Bodies ; the Real Distances and Dimensions ; Universal Gravitation. The most recent Astronomical Discoveries are incorporated. Mr. Lockyer's work supplements that of the Astronomer Royal. "The book is full, clear, sound, and worthy of attention, not only as a popular exposition, but as a scientific 'Index.'—Athenæum. "The most fascinating of elementary books on the Sciences."—Nonconformist.

Macmillan (Rev. Hugh).—For other Works by the same Author, see THEOLOGICAL CATALOGUE.

HOLIDAYS ON HIGH LANDS ; or, Rambles and Incidents in search of Alpine Plants. Crown 8vo. cloth. 6s.

The aim of this book is to impart a general idea of the origin, character, and distribution of those rare and beautiful Alpine plants which occur on the British hills, and which are found almost everywhere on the lofty mountain chains of Europe, Asia, Africa, and America. In the first three chapters the peculiar vegetation of the Highland mountains is fully described ; while in the remaining chapters this vegetation is traced to its northern cradle in the mountains of Norway, and to its southern European termination in the Alps of Switzerland. The information the author has to give is conveyed in a setting of personal adventure. "One of the most charming books of its kind ever written."—Literary Churchman. "Mr. M.'s glowing pictures of Scandinavian scenery."—Saturday Review.

FOOT-NOTES FROM THE PAGE OF NATURE. With numerous Illustrations. Fcap. 8vo. 5s.

"Those who have derived pleasure and profit from the study of flowers and ferns—subjects, it is pleasing to find, now everywhere popular—by descending lower into the arcana of the vegetable kingdom, will find a still more interesting and delightful field of research in the objects brought under review in the following pages."—Preface. "The naturalist and the botanist will delight in this volume, and those who understand little of the scientific parts of the work will linger over the mysterious page of nature here unfolded to their view."—John Bull.

Mansfield (C. B.)—A THEORY OF SALTS. A Treatise on the Constitution of Bipolar (two-membered) Chemical Compounds. By the late CHARLES BLACHFORD MANSFIELD. Crown 8vo. 14s.

"Mansfield," says the editor, "wrote this book to defend the principle that the fact of voltaic decomposition afforded the true indication, if properly interpreted, of the nature of the saline structure, and of the atomicity of the elements that built it up. No chemist will peruse this book without feeling that he is in the presence of an

original thinker, whose pages are continually suggestive, even though their general argument may not be entirely concurrent in direction with that of modern chemical thought."

Mivart (St. George).—ON THE GENESIS OF SPECIES.

By ST. GEORGE MIVART, F.R.S. Crown 8vo. Second Edition, to which notes have been added in reference and reply to Darwin's "Descent of Man." With numerous Illustrations. pp. xv. 296. 9s.

The aim of the author is to support the doctrine that the various species have been evolved by ordinary natural laws (for the most part unknown) controlled by the subordinate action of "natural selection," and at the same time to remind some that there is and can be absolutely nothing in physical science which forbids them to regard those natural laws as acting with the Divine concurrence, and in obedience to a creative fiat originally imposed on the primeval cosmos, "in the beginning," by its Creator, its Upholder, and its Lord. Nearly fifty woodcuts illustrate the letter-press, and a complete index makes all references extremely easy. Canon Kingsley, in his address to the "Devonshire Association," says, "Let me recommend earnestly to you, as a specimen of what can be said on the other side, the 'Genesis of Species,' by Mr. St. George Mivart, F.R.S., a book which I am happy to say has been received elsewhere as it has deserved, and, I trust, will be received so among you." "In no work in the English language has this great controversy been treated at once with the same broad and vigorous grasp of facts, and the same liberal and candid temper."—Saturday Review.

Nature.—A WEEKLY ILLUSTRATED JOURNAL OF SCIENCE. Published every Thursday. Price 4d. Monthly Parts, 1s. 4d. and 1s. 8d.; Half-yearly Volumes, 10s. 6d. Cases for binding vols. 1s. 6d.

"Backed by many of the best names among English philosophers, and by a few equally valuable supporters in America and on the Continent of Europe."—Saturday Review. "This able and well-edited Journal, which posts up the science of the day promptly, and promises to be of signal service to students and savants."—British Quarterly Review.

Oliver.—Works by DANIEL OLIVER, F.R.S., F.L.S., Professor of Botany in University College, London, and Keeper of the Herbarium and Library of the Royal Gardens, Kew :—

LESSONS IN ELEMENTARY BOTANY. With nearly Two Hundred Illustrations. Twelfth Thousand. 18mo cloth. 4s. 6d.

This book is designed to teach the elements of Botany on Professor Henslow's plan of selected Types and by the use of Schedules. The earlier chapters, embracing the elements of Structural and Physiological Botany, introduce us to the methodical study of the Ordinal Types. The concluding chapters are entitled, "How to Dry Plants" and "How to Describe Plants." A valuable Glossary is appended to the volume. In the preparation of this work free use has been made of the manuscript materials of the late Professor Henslow.

FIRST BOOK OF INDIAN BOTANY. With numerous Illustrations. Extra fcap. 8vo. 6s. 6d.

This manual is, in substance, the author's "Lessons in Elementary Botany," adapted for use in India. In preparing it he has had in view the want, often felt, of some handy résumé of Indian Botany, which might be serviceable not only to residents of India, but also to any one about to proceed thither, desirous of getting some preliminary idea of the botany of the country. It contains a well-digested summary of all essential knowledge pertaining to Indian Botany, wrought out in accordance with the best principles of scientific arrangement."—Allen's Indian Mail.

Penrose (F. C.)—ON A METHOD OF PREDICTING BY GRAPHICAL CONSTRUCTION, OCCULTATIONS OF STARS BY THE MOON, AND SOLAR ECLIPSES FOR ANY GIVEN PLACE. Together with more rigorous methods for the Accurate Calculation of Longitude. By F. C. PENROSE, F.R.A.S. With Charts, Tables, etc. 4to. 12s.

The author believes that if, by a graphic method, the prediction of occultations can be rendered more inviting, as well as more expeditious, than by the method of calculation, it may prove acceptable to the nautical profession as well as to scientific travellers or amateurs. The author has endeavoured to make the whole process as intelligible as possible, so that the beginner, instead of merely having to

follow directions imperfectly understood, may readily comprehend the meaning of each step, and be able to illustrate the practice by the theory. Besides all necessary charts and tables, the work contains a large number of skeleton forms for working out cases in practice.

Roscoe.—Works by HENRY E. ROSCOE, F.R.S., Professor of Chemistry in Owens College, Manchester :—

LESSONS IN ELEMENTARY CHEMISTRY, INORGANIC AND ORGANIC. With numerous Illustrations and Chromolitho of the Solar Spectrum, and of the Alkalies and Alkaline Earths. New Edition. Thirty-first Thousand. 18mo. cloth. 4s. 6d.

It has been the endeavour of the author to arrange the most important facts and principles of Modern Chemistry in a plain but concise and scientific form, suited to the present requirements of elementary instruction. For the purpose of facilitating the attainment of exactitude in the knowledge of the subject, a series of exercises and questions upon the lessons have been added. The metric system of weights and measures, and the centigrade thermometric scale, are used throughout this work. The new edition, besides new woodcuts, contains many additions and improvements, and includes the most important of the latest discoveries. "We unhesitatingly pronounce it the best of all our elementary treatises on Chemistry."—Medical Times.

SPECTRUM ANALYSIS. Six Lectures, with Appendices, Engravings, Maps, and Chromolithographs. Royal 8vo. 21s.

A Second Edition of these popular Lectures, containing all the most recent discoveries and several additional illustrations. "In six lectures he has given the history of the discovery and set forth the facts relating to the analysis of light in such a way that any reader of ordinary intelligence and information will be able to understand what 'Spectrum Analysis' is, and what are its claims to rank among the most signal triumphs of science."—Nonconformist. "The lectures themselves furnish a most admirable elementary treatise on the subject, whilst by the insertion in appendices to each lecture of extracts from the most important published memoirs, the author has rendered it equally valuable as a text-book for advanced students."—Westminster Review.

Stewart (B.)—LESSONS IN ELEMENTARY PHYSICS.

By BALFOUR STEWART, F.R.S., Professor of Natural Philosophy in Owens College, Manchester. With numerous Illustrations and Chromolithos of the Spectra of the Sun, Stars, and Nebulæ. Second Edition. 18mo. 4s. 6d.

A description, in an elementary manner, of the most important of those laws which regulate the phenomena of nature. The active agents, heat, light, electricity, etc., are regarded as varieties of energy, and the work is so arranged that their relation to one another, looked at in this light, and the paramount importance of the laws of energy, are clearly brought out. The volume contains all the necessary illustrations. The Educational Times calls this "the bean-ideal of a scientific text-book, clear, accurate, and thorough."

Thudichum and Dupré.—A TREATISE ON THE ORIGIN, NATURE, AND VARIETIES OF WINE.

Being a Complete Manual of Viticulture and Oenology. By J. L. W. THUDICHUM, M.D., and AUGUST DUPRÉ, Ph.D., Lecturer on Chemistry at Westminster Hospital. Medium 8vo. cloth gilt. 25s.

In this elaborate work the subject of the manufacture of wine is treated scientifically in minute detail, from every point of view. A chapter is devoted to the Origin and Physiology of Vines, two to the Principles of Viticulture; while other chapters treat of Vintage and Vinification, the Chemistry of Alcohol, the Acids, Ether, Sugars, and other matters occurring in wine. This introductory matter occupies the first nine chapters, the remaining seventeen chapters being occupied with a detailed account of the Viticulture and the Wines of the various countries of Europe, of the Atlantic Islands, of Asia, of Africa, of America, and of Australia. Besides a number of Analytical and Statistical Tables, the work is enriched with eighty-five illustrative woodcuts. "A treatise almost unique for its usefulness either to the wine-grower, the vendor, or the consumer of wine. The analyses of wine are the most complete we have yet seen, exhibiting at a glance the constituent principles of nearly all the wines known in this country."—Wine Trade Review.

Wallace (A. R.)—CONTRIBUTIONS TO THE THEORY OF NATURAL SELECTION. A Series of Essays. By

ALFRED RUSSEL WALLACE, Author of "The Malay Archipelago," etc. Second Edition, with Corrections and Additions. Crown 8vo. 8s.6d. (For other Works by the same Author, see CATALOGUE OF HISTORY AND TRAVELS.)

Mr. Wallace has good claims to be considered as an independent originator of the theory of natural selection. Dr. Hooker, in his address to the British Association, spoke thus of the author: "Of Mr. Wallace and his many contributions to philosophical biology it is not easy to speak without enthusiasm; for, putting aside their great merits, he, throughout his writings, with a modesty as rare as I believe it to be unconscious, forgets his own unquestioned claim to the honour of having originated, independently of Mr. Darwin, the theories which he so ably defends." The Saturday Review says: "He has combined an abundance of fresh and original facts with a liveliness and sagacity of reasoning which are not often displayed so effectively on so small a scale." The Essays in this volume are:—I. "On the Law which has regulated the introduction of New Species." II. "On the Tendencies of Varieties to depart indefinitely from the Original Type." III. "Mimicry, and other Protective Resemblances among Animals." IV. "The Malayan Papilionida, as illustrative of the Theory of Natural Selection." V. "On Instinct in Man and Animals." VI. "The Philosophy of Birds' Nests." VII. "A Theory of Birds' Nests." VIII. "Creation by Law." IX. "The Development of Human Races under the Law of Natural Selection." X. "The Limits of Natural Selection as applied to Man."

Warrington.—THE WEEK OF CREATION; OR, THE COSMOGONY OF GENESIS CONSIDERED IN ITS RELATION TO MODERN SCIENCE. By GEORGE WARRINGTON, Author of "The Historic Character of the Pentateuch Vindicated." Crown 8vo. 4s. 6d.

The greater part of this work is taken up with the teaching of the Cosmogony. Its purpose is also investigated, and a chapter is devoted to the consideration of the passage in which the difficulties occur. "A very able vindication of the Mosaic Cosmogony, by a writer who unites the advantages of a critical knowledge of the Hebrew text and of distinguished scientific attainments."—Spectator.

Wilson.—Works by the late GEORGE WILSON, M.D., F.R.S.E.,
Regius Professor of Technology in the University of Edinburgh :—

RELIGIO CHEMICI. With a Vignette beautifully engraved after
a design by Sir NOEL PATON. Crown 8vo. 8s. 6d.

"George Wilson," says the Preface to this volume, "had it in his heart for many years to write a book corresponding to the *Religio Medici* of Sir Thomas Browne, with the title *Religio Chemicæ*. Several of the Essays in this volume were intended to form chapters of it. These fragments being in most cases like finished gems waiting to be set, some of them are now given in a collected form to his friends and the public. In living remembrance of his purpose, the name chosen by himself has been adopted, although the original design can be but very faintly represented." The Contents of the volume are:—"Chemistry and Natural Theology." "The Chemistry of the Stars; an Argument touching the Stars and their Inhabitants." "Chemical Final Causes; as illustrated by the presence of Phosphorus, Nitrogen, and Iron in the Higher Sentient Organisms." "Robert Boyle." "Wollaston." "Life and Discoveries of Dalton." "Thoughts on the Resurrection; an Address to Medical Students." "A more fascinating volume," the Spectator says, "has seldom fallen into our hands." The Freeman says: "These papers are all valuable and deeply interesting. The production of a profound thinker, a suggestive and eloquent writer, and a man whose piety and genius went hand in hand."

THE PROGRESS OF THE TELEGRAPH. Fcap. 8vo. 1s.

"While a complete view of the progress of the greatest of human inventions is obtained, all its suggestions are brought out with a rare thoughtfulness, a genial humour, and an exceeding beauty of utterance."—Nonconformist.

Winslow.—**FORCE AND NATURE: ATTRACTION AND REPULSION.** The Radical Principles of Energy graphically discussed in their Relations to Physical and Morphological Development. By C. F. WINSLOW, M.D. 8vo. 14s.

The author having for long investigated Nature in many directions, has ever felt unsatisfied with the physical foundations upon which some branches of science have been so long compelled to rest. The question, he believes, must have occurred to many astronomers and

physicists whether some subtle principle antagonistic to attraction does not also exist as an all-pervading element in nature, and so operate as in some way to disturb the action of what is generally considered by the scientific world a unique force. The aim of the present work is to set forth this subject in its broadest aspects, and in such a manner as to invite thereto the attention of the learned. The subjects of the eleven chapters are:—I. "Space." II. "Matter." III. "Inertia, Force, and Mind." IV. "Molecules." V. "Molecular Force." VI. "Union and Inseparability of Matter and Force." VII. and VIII. "Nature and Action of Force—Attraction—Repulsion." IX. "Cosmical Repulsion. X. "Mechanical Force." XI. "Central Forces and Celestial Physics." "Deserves thoughtful and conscientious study."—Saturday Review.

Wurtz.—A HISTORY OF CHEMICAL THEORY, from the Age of Lavoisier down to the present time. By AD. WURTZ. Translated by HENRY WATTS, F.R.S. Crown 8vo. 6s.

"The discourse, as a résumé of chemical theory and research, unites singular luminousness and grasp. A few judicious notes are added by the translator."—Pall Mall Gazette. "The treatment of the subject is admirable, and the translator has evidently done his duty most efficiently."—Westminster Review.

WORKS IN PHYSIOLOGY, ANATOMY, AND MEDICAL WORKS GENERALLY.

Allbutt (T. C.)—ON THE USE OF THE OPHTHALMOSCOPE in Diseases of the Nervous System and of the Kidneys; also in certain other General Disorders. By THOMAS CLIFFORD ALLBUTT, M.A., M.D. Cantab., Physician to the Leeds General Infirmary, Lecturer on Practical Medicine, etc. etc. 8vo. 15s.

The Ophthalmoscope has been found of the highest value in the investigation of nervous diseases. But it is not easy for physicians who have left the schools, and are engaged in practice, to take up a new

instrument which requires much skill in using ; it is therefore hoped that by such the present volume, containing the results of the author's extensive use of the instrument in diseases of the nervous system, will be found of high value ; and that to all students it may prove a useful hand-book. After four introductory chapters on the history and value of the Ophthalmoscope, and the manner of investigating the states of the optic nerve and retina, the author treats of the various diseases with which optic changes are associated, and describes the way in which such associations take place. Besides the cases referred to throughout the volume, the Appendix contains details of 123 cases illustrative of the subjects discussed in the text, and a series of tabulated cases to show the Ophthalmoscopic appearances of the eye in Insanity, Mania, Dementia, Melancholia and Monomania, Idiocy, and General Paralysis. The volume is illustrated with two valuable coloured plates of morbid appearances of the eye under the Ophthalmoscope. "By its aid men will no longer be compelled to work for years in the dark ; they will have a definite standpoint whence to proceed on their course of investigation." —Medical Times.

Anstie (F. E.)—NEURALGIA, AND DISEASES WHICH RESEMBLE IT. By FRANCIS E. ANSTIE, M.D., M.R.C.P., Senior Assistant Physician to Westminster Hospital. 8vo. 10s. 6d.

Dr. Anstie is well known as one of the greatest living authorities on Neuralgia. The present treatise is the result of many years' careful independent scientific investigation into the nature and proper treatment of this most painful disease. The author has had abundant means of studying the subject both in his own person and in the hundreds of patients that have resorted to him for treatment. He has gone into the whole subject indicated in the title ab initio, and the publishers believe it will be found that he has presented it in an entirely original light, and done much to rob this excruciating and hitherto refractory disease of many of its terrors. The Introduction treats briefly of Pain in General, and contains some striking and even original ideas as to its nature and in reference to sensation generally.

Barwell.—THE CAUSES AND TREATMENT OF LATERAL CURVATURE OF THE SPINE. Enlarged from Lectures published in the *Lancet*. By RICHARD BARWELL, F.R.C.S.,

Surgeon to and Lecturer on Anatomy at the Charing Cross Hospital.
Second Edition. Crown 8vo. 4s. 6d.

Having failed to find in books a satisfactory theory of those conditions which produce lateral curvature, Mr. Barwell resolved to investigate the subject for himself ab initio. The present work is the result of long and patient study of Spines, normal and abnormal. He believes the views which he has been led to form account for those essential characteristics which have hitherto been left unexplained; and the treatment which he advocates is certainly less irksome, and will be found more efficacious than that which has hitherto been pursued. Indeed, the mode in which the first edition has been received by the profession is a gratifying sign that Mr. Barwell's principles have made their value and their weight felt. Many pages and a number of woodcuts have been added to the Second Edition.

Corfield (Professor W. H.)—A DIGEST OF FACTS RELATING TO THE TREATMENT AND UTILIZATION OF SEWAGE. By W. H. CORFIELD, M.A., B.A., Professor of Hygiene and Public Health at University College, London. 8vo. 10s. 6d. Second Edition, corrected and enlarged.

The author in the Second Edition has revised and corrected the entire work, and made many important additions. The headings of the eleven chapters are as follow:—I. "Early Systems: Midden-Heaps and Cesspools." II. "Filth and Disease—Cause and Effect." III. "Improved Midden-Pits and Cesspools; Midden-Closets, Pail-Closets, etc." IV. "The Dry-Closet Systems. V. "Water-Closets." VI. "Sewerage." VII. "Sanitary Aspects of the Water-Carrying System." VIII. "Value of Sewage; Injury to Rivers." IX. "Town Sewage; Attempts at Utilization." X. "Filtration and Irrigation." XI. "Influence of Sewage Farming on the Public Health." An abridged account of the more recently published researches on the subject will be found in the Appendices, while the Summary contains a concise statement of the views which the author himself has been led to adopt: references have been inserted throughout to show from what sources the numerous quotations have been derived, and an Index has been added. "Mr. Corfield's work is entitled to rank as a standard authority, no less than a convenient handbook, in all matters relating to sewage."—Athenæum.

Elam (C.)—A PHYSICIAN'S PROBLEMS. By CHARLES ELAM, M.D., M.R.C.P. Crown 8vo. 9s.

CONTENTS :—“*Natural Heritage.*” “*On Degeneration in Man.*” “*On Moral and Criminal Epidemics.*” “*Body v. Mind.*” “*Illusions and Hallucinations.*” “*On Somnambulism.*” “*Reverie and Abstraction.*” These Essays are intended as a contribution to the *Natural History* of those outlying regions of *Thought and Action* whose domain is the debateable ground of *Brain, Nerve, and Mind.* They are designed also to indicate the origin and mode of perpetuation of those varieties of organization, intelligence, and general tendencies towards *vice or virtue*, which seem to be so capriciously developed among mankind. They also point to causes for the infinitely varied forms of disorder of nerve and brain—organic and functional—far deeper and more recondite than those generally believed in. “*The book is one which all statesmen, magistrates, clergymen, medical men, and parents should study and inwardly digest.*”—Examiner.

Fox.—Works by WILSON FOX, M.D. Lond., F.R.C.P., Holme Professor of Clinical Medicine, University College, London, Physician Extraordinary to her Majesty the Queen, etc. :—

ON THE DIAGNOSIS AND TREATMENT OF THE VARIETIES OF DYSPEPSIA, CONSIDERED IN RELATION TO THE PATHOLOGICAL ORIGIN OF THE DIFFERENT FORMS OF INDIGESTION. Second Edition. 8vo. 7s. 6d.

ON THE ARTIFICIAL PRODUCTION OF TUBERCLE IN THE LOWER ANIMALS, With Coloured Plates. 4to. 5s. 6d.

In this Lecture Dr. Fox describes in minute detail a large number of experiments made by him on guinea-pigs and rabbits for the purpose of inquiring into the origin of Tubercle by the agency of direct irritation or by septic matters. This method of inquiry he believes to be one of the most important advances which have been recently made in the pathology of the disease. The work is illustrated by three plates, each containing a number of carefully coloured illustrations from nature.

ON THE TREATMENT OF HYPERPYREXIA, as Illustrated in Acute Articular Rheumatism by means of the External Application of Cold. 8vo. 2s. 6d.

The object of this work is to show that the class of cases included under the title, and which have hitherto been invariably fatal, may, by a judicious use of the cold bath and without venesection, be brought to a favourable termination. Minute details are given of the successful treatment by this method of two patients by the author, followed by a Commentary on the cases, in which the merits of the mode of treatment are discussed and compared with those of methods followed by other eminent practitioners. Appended are tables of the observations made on the temperature during the treatment; a table showing the effect of the immersion of the patients in the baths employed, in order to exhibit the rate at which the temperature was lowered in each case; a table of the chief details of twenty-two cases of this class recently published, and which are referred to in various parts of the Commentary. Two Charts are also introduced, giving a connected view of the progress of the two successful cases, and a series of sphygmographic tracings of the pulses of the two patients. "A clinical study of rare value. Should be read by every one."—Medical Press and Circular.

Galton (D.)—AN ADDRESS ON THE GENERAL PRINCIPLES WHICH SHOULD BE OBSERVED IN THE CONSTRUCTION OF HOSPITALS. Delivered to the British Medical Association at Leeds, July 1869. By DOUGLAS GALTON, C.B., F.R.S. Crown 8vo. 3s. 6d.

In this Address the author endeavours to enunciate what are those principles which seem to him to form the starting-point from which all architects should proceed in the construction of hospitals. Besides Mr. Galton's paper the book contains the opinions expressed in the subsequent discussion by several eminent medical men, such as Dr. Kennedy, Sir James Y. Simpson, Dr. Hughes Bennet, and others. The work is illustrated by a number of plans, sections, and other cuts. "An admirable exposition of those conditions of structure which most conduce to cleanliness, economy, and convenience."—Times.

Harley (J.)—THE OLD VEGETABLE NEUROTICS, Hemlock, Opium, Belladonna, and Henbane; their Physiological Action and Therapeutical Use, alone and in combination. Being the Gulstonian Lectures of 1868 extended, and including a Complete Examination of the Active Constituents of Opium. By JOHN HARLEY, M.D. Lond., F.R.C.P., F.L.S., etc. 8vo. 12s.

The author's object throughout the investigations and experiments on which this volume is founded has been to ascertain, clearly and definitely, the action of the drugs employed on the healthy body in medicinal doses, from the smallest to the largest; to deduce simple practical conclusions from the facts observed; and then to apply the drug to the relief of the particular conditions to which its action appeared suited. Many experiments have been made by the author both on men and the lower animals; and the author's endeavour has been to present to the mind, as far as words may do, impressions of the actual condition of the individual subjected to the drug. "Those who are interested generally in the progress of medical science will find much to repay a careful perusal."—Athenæum.

Hood (Wharton).—ON BONE-SETTING (so called), and its Relation to the Treatment of Joints Crippled by Injury, Rheumatism, Inflammation, etc. etc. By WHARTON P. HOOD, M.D., M.R.C.S. Crown 8vo. 4s. 6d.

The author for a period attended the London practice of the late Mr. Hutton, the famous and successful bone-setter, by whom he was initiated into the mystery of the art and practice. Thus the author is amply qualified to write on the subject from the practical point of view, while his professional education enables him to consider it in its scientific and surgical bearings. In the present work he gives a brief account of the salient features of a bone-setter's method of procedure in the treatment of damaged joints, of the results of that treatment, and of the class of cases in which he has seen it prove successful. The author's aim is to give the rationale of the bone-setter's practice, to reduce it to something like a scientific method, to show when force should be resorted to and when it should not, and to initiate surgeons into the secret of Mr. Hutton's successful manipulation. Throughout the work a great number of authentic instances of successful treatment are given, with the details of the method of cure; and the Chapters on Manipulations and Affections of the Spine are illustrated by a number of appropriate and well-executed cuts. "Dr. Hood's book is full of instruction, and should be read by all surgeons."—Medical Times.

Humphry.—THE HUMAN SKELETON (including the joints). By G. M. HUMPHRY, M.D., F.R.S. With 260 Illustrations, drawn from nature. Medium 8vo. 28s.

In lecturing on the Skeleton it has been the author's practice, instead of giving a detailed account of the several parts, to request his students to get up the descriptive anatomy of certain bones, with the aid of some work on osteology. He afterwards tested their acquirements by examination, endeavouring to supply deficiencies and correct errors, adding also such information—physical, physiological, pathological, and practical—as he had gathered from his own observation and researches, and which was likely to be useful and excite an interest in the subject. This additional information forms, in great part, the material of this volume, which is intended to be supplementary to existing works on anatomy. Considerable space has been devoted to the description of the joints, because it is less fully given in other works, and because an accurate knowledge of the structure and peculiar form of the joints is essential to a correct knowledge of their movements. The numerous illustrations were all drawn upon stone from nature; and in most instances, from specimens prepared for the purpose by the author himself. "Bearing at once the stamp of the accomplished scholar, and evidences of the skilful anatomist. We express our admiration of the drawings."—Medical Times and Gazette.

Huxley's Physiology.—See p. 24, preceding.

Journal of Anatomy and Physiology.

Conducted by Professors HUMPHRY and NEWTON, and Mr. CLARK of Cambridge, Professor TURNER of Edinburgh, and Dr. WRIGHT of Dublin. Published twice a year. Old Series, Parts I. and II., price 7s. 6d. each. Vol. I. containing Parts I. and II., Royal 8vo., 16s. New Series, Parts I. to IX. 6s. each, or yearly Vols. 12s. 6d. each.

Lankester.—COMPARATIVE LONGEVITY IN MAN AND THE LOWER ANIMALS. By E. RAY LANKESTER, B.A. Crown 8vo. 4s. 6d.

This Essay gained the prize offered by the University of Oxford for the best Paper on the subject of which it treats. This interesting subject is here treated in a thorough manner, both scientifically and statistically.

Maclaren.—TRAINING, IN THEORY AND PRACTICE. By ARCHIBALD MACLAREN, the Gymnasium, Oxford. 8vo. Handsomely bound in cloth, 7s. 6d.

The ordinary agents of health are Exercise, Diet, Sleep, Air, Bathing, and Clothing. In this work the author examines each of these agents in detail, and from two different points of view. First, as to the manner in which it is, or should be, administered under ordinary circumstances : and secondly, in what manner and to what extent this mode of administration is, or should be, altered for purposes of training ; the object of "training," according to the author, being "to put the body, with extreme and exceptional care, under the influence of all the agents which promote its health and strength, in order to enable it to meet extreme and exceptional demands upon its energies." Appended are various diagrams and tables relating to boat-racing, and tables connected with diet and training. "The philosophy of human health has seldom received so apt an exposition."—Globe. "After all the nonsense that has been written about training, it is a comfort to get hold of a thoroughly sensible book at last."—John Bull.

Macpherson.—Works by JOHN MACPHERSON, M.D. :—

THE BATHS AND WELLS OF EUROPE ; Their Action and Uses. With Hints on Change of Air and Diet Cures. With a Map. Extra fcap. 8vo. 6s. 6d.

This work is intended to supply information which will afford aid in the selection of such Spas as are suited for particular cases. It exhibits a sketch of the present condition of our knowledge on the subject of the operation of mineral waters, gathered from the author's personal observation, and from every other available source of information. It is divided into four books, and each book into several chapters :—Book I. Elements of Treatment, in which, among other matters, the external and internal uses of water are treated of. II. Bathing, treating of the various kinds of baths. III. Wells, treating of the various kinds of mineral waters. IV. Diet Cures, in which various vegetable, milk, and other "cures" are discussed. Appended is an Index of Diseases noticed, and one of places named. Prefixed is a sketch map of the principal baths and places of health-resort in Europe. "Dr. Macpherson has given the kind of information which every medical practitioner ought to possess."—The Lancet. "Whoever wants to know the real character of any health-resort must read Dr. Macpherson's book."—Medical Times.

Macpherson (J.)—*continued.*

OUR BATHS AND WELLS : The Mineral Waters of the British Islands, with a List of Sea-bathing Places. Extra fcap. 8vo. pp. xv. 205. 3s. 6d.

Dr. Macpherson has divided his work into five parts. He begins by a few introductory observations on bath life, its circumstances, uses, and pleasures; he then explains in detail the composition of the various mineral waters, and points out the special curative properties of each class. A chapter on "The History of British Wells" from the earliest period to the present time forms the natural transition to the second part of this volume, which treats of the different kinds of mineral waters in England, whether pure, thermal and earthy, saline, chalybeate, or sulphur. Wales, Scotland, and Ireland supply the materials for distinct sections. An Index of mineral waters, one of sea-bathing places, and a third of wells of pure or nearly pure water, terminate the book. "This little volume forms a very available handbook for a large class of invalids."—Nonconformist.

Maudsley.—Works by HENRY MAUDSLEY, M.D., Professor of Medical Jurisprudence in University College, London :—

BODY AND MIND : An Inquiry into their Connection and Mutual Influence, specially in reference to Mental Disorders; being the Gulstonian Lectures for 1870. Delivered before the Royal College of Physicians. Crown 8vo. 5s.

The volume consists of three Lectures and two long Appendices, the general plan of the whole being to bring Man, both in his physical and mental relations, as much as possible under the scope of scientific inquiry. The first Lecture is devoted to an exposition of the physical conditions of mental function in health. In the second Lecture are sketched the features of some forms of degeneracy of mind, as exhibited in morbid varieties of the human kind, with the purpose of bringing prominently into notice the operation of physical causes from generation to generation, and the relationship of mental to other diseases of the nervous system. In the third Lecture are displayed the relations of morbid states of the body and disordered mental function. Appendix I. is a criticism of the Archbishop of York's address on "The Limits of Philosophical Inquiry." Appendix II. deals with the "Theory of Vitality," in which the author en-

Maudsley (H.)—*continued.*

deavours to set forth the reflections which facts seem to warrant.
"It distinctly marks a step in the progress of scientific psychology."
 —The Practitioner.

THE PHYSIOLOGY AND PATHOLOGY OF MIND.
 Second Edition, Revised. 8vo. 16s.

This work is the result of an endeavour on the author's part to arrive at some definite conviction with regard to the physical conditions of mental function, and the relation of the phenomena of sound and unsound mind. The author's aim throughout has been twofold : I. To treat of mental phenomena from a physiological rather than from a metaphysical point of view. II. To bring the manifold instructive instances presented by the unsound mind to bear upon the interpretation of the obscure problems of mental science. In the first part, the author pursues his independent inquiry into the science of Mind in the same direction as that followed by Bain, Spencer, Laycock, and Carpenter ; and in the second, he studies the subject in a light which, in this country at least, is almost entirely novel: "Dr. Maudsley's work, which has already become standard, we most urgently recommend to the careful study of all those who are interested in the physiology and pathology of the brain."—Anthropological Review.

Practitioner (The).—A Monthly Journal of Therapeutics.
 Edited by FRANCIS E. ANSTIE, M.D. 8vo. Price 1s. 6d.
 Vols. I to VII. 8vo. cloth. 10s. 6d. each.**Radcliffe.**—DYNAMICS OF NERVE AND MUSCLE. By CHARLES BLAND RADCLIFFE, M.D., F.R.C.P., Physician to the Westminster Hospital, and to the National Hospital for the Paralysed and Epileptic. Crown 8vo. 8s. 6d.

This work contains the result of the author's long investigations into the Dynamics of Nerve and Muscle, as connected with Animal Electricity. The author endeavours to show from these researches that the state of action in nerve and muscle, instead of being a manifestation of vitality, must be brought under the domain of physical law in order to be intelligible, and that a different meaning, also based upon pure physics, must be attached to the state of rest. "The practitioner

will find in Dr. Radcliffe a 'guide, philosopher, and friend,' from whose teaching he cannot fail to reap a plentiful harvest of new and valuable ideas."—Scotsman.

Reynolds.—A SYSTEM OF MEDICINE. Vol. I. Edited by J. RUSSELL REYNOLDS, M.D., F.R.C.P. London. Second Edition. 8vo. 25s.

Part I. General Diseases, or Affections of the Whole System.

§ I.—*Those determined by agents operating from without, such as the exanthemata, malarial diseases, and their allies.* § II.—*Those determined by conditions existing within the body, such as Gout, Rheumatism, Rickets, etc.* *Part II. Local Diseases, or Affections of particular Systems.* § I.—*Diseases of the Skin.*

A SYSTEM OF MEDICINE. Vol. II. Second Edition in the Press. 8vo. 25s.

Part II. Local Diseases (continued). § I.—*Diseases of the Nervous System.* A. *General Nervous Diseases.* B. *Partial Diseases of the Nervous System.* 1. *Diseases of the Head.* 2. *Diseases of the Spinal Column.* 3. *Diseases of the Nerves.* § II.—*Diseases of the Digestive System.* A. *Diseases of the Stomach.*

A SYSTEM OF MEDICINE. Vol. III. 8vo. 25s.

Part II. Local Diseases (continued). § II. *Diseases of the Digestive System (continued).* B. *Diseases of the Mouth.* C. *Diseases of the Fauces, Pharynx, and Œsophagus.* D. *Diseases of the Intestines.* E. *Diseases of the Peritoneum.* F. *Diseases of the Liver.* G. *Diseases of the Pancreas.* § III.—*Diseases of the Respiratory System.* A. *Diseases of the Larynx.* B. *Diseases of the Thoracic Organs.* "One of the best and most comprehensive treatises on Medicine which have yet been attempted in any country."—*Indian Medical Journal.* "Contains some of the best essays that have lately appeared, and is a complete library in itself."—*Medical Press.*

Seaton.—A HANDBOOK OF VACCINATION. By EDWARD C. SEATON, M.D., Medical Inspector to the Privy Council. Extra fcap. 8vo. 8s. 6d.

The author's object in putting forth this work is twofold: First, to provide a text-book on the science and practice of Vaccination for

the use of younger practitioners and of medical students ; secondly, to give what assistance he could to those engaged in the administration of the system of Public Vaccination established in England. For many years past, from the nature of his office, Dr. Seaton has had constant intercourse in reference to the subject of Vaccination, with medical men who are interested in it, and especially with that large part of the profession who are engaged as Public Vaccinators. All the varieties of pocks, both in men and the lower animals, are treated of in detail, and much valuable information given on all points connected with lymph, and minute instructions as to the niceties and cautions which so greatly influence success in Vaccination. The administrative sections of the work will be of interest and value, not only to medical practitioners, but to many others to whom a right understanding of the principles on which a system of Public Vaccination should be based is indispensable. "Henceforth the indispensable handbook of Public Vaccination, and the standard authority on this great subject."—British Medical Journal.

Symonds (J. A., M.D.)—MISCELLANIES. By JOHN ADDINGTON SYMONDS, M.D. Selected and Edited, with an Introductory Memoir, by his Son. 8vo. 7s. 6d.

The late Dr. Symonds of Bristol was a man of a singularly versatile and elegant as well as powerful and scientific intellect. In order to make this selection from his many works generally interesting, the editor has confined himself to works of pure literature, and to such scientific studies as had a general philosophical or social interest. Among the general subjects are articles on "the Principles of Beauty," on "Knowledge," and a "Life of Dr. Prichard ;" among the Scientific Studies are papers on "Sleep and Dreams," "Apparitions," "the Relations between Mind and Muscle," "Habit," etc. ; there are several papers on "the Social and Political Aspects of Medicine ;" and a few Poems and Translations selected from a great number of equal merit. "A collection of graceful essays on general and scientific subjects, by a very accomplished physician."—Graphic.

WORKS ON MENTAL AND MORAL PHILOSOPHY, AND ALLIED SUBJECTS.

Aristotle.—AN INTRODUCTION TO ARISTOTLE'S RHETORIC. With Analysis, Notes, and Appendices. By E. M. COPE, Trinity College, Cambridge. 8vo. 14s.

This work is introductory to an edition of the Greek Text of Aristotle's Rhetoric, which is in course of preparation. Its object is to render that treatise thoroughly intelligible. The author has aimed to illustrate, as preparatory to the detailed explanation of the work, the general bearings and relations of the Art of Rhetoric in itself, as well as the special mode of treating it adopted by Aristotle in his peculiar system. The evidence upon obscure or doubtful questions connected with the subject is examined; and the relations which Rhetoric bears, in Aristotle's view, to the kindred art of Logic are fully considered. A connected Analysis of the work is given, and a few important matters are separately discussed in Appendices. There is added, as a general Appendix, by way of specimen of the antagonistic system of Isocrates and others, a complete analysis of the treatise called Ῥητορικὴ πρὸς Ἀλέξανδρον, with a discussion of its authorship and of the probable results of its teaching.

ARISTOTLE ON FALLACIES ; OR, THE SOPHISTICI ELENCHI. With a Translation and Notes by EDWARD POSTE, M A., Fellow of Oriel College, Oxford. 8vo. 8s. 6d.

Besides the doctrine of Fallacies, Aristotle offers, either in this treatise or in other passages quoted in the Commentary, various glances over the world of science and opinion, various suggestions or problems which are still agitated, and a vivid picture of the ancient system of dialectics, which it is hoped may be found both interesting

and instructive. "It will be an assistance to genuine students of Aristotle."—Guardian. "It is indeed a work of great skill."—Saturday Review.

Butler (W. A.), Late Professor of Moral Philosophy in the University of Dublin :—

LECTURES ON THE HISTORY OF ANCIENT PHILOSOPHY. Edited from the Author's MSS., with Notes, by WILLIAM HEPWORTH THOMPSON, M.A., Master of Trinity College, and Regius Professor of Greek in the University of Cambridge. Two Volumes. 8vo. 1l. 5s.

These Lectures consist of an Introductory Series on the Science of Mind generally, and five other Series on Ancient Philosophy, the greater part of which treat of Plato and the Platonists, the Fifth Series being an unfinished course on the Psychology of Aristotle, containing an able Analysis of the well known though by no means well understood Treatise, περὶ ψυχῆς. These Lectures are the result of patient and conscientious examination of the original documents, and may be considered as a perfectly independent contribution to our knowledge of the great master of Grecian wisdom. The author's intimate familiarity with the metaphysical writings of the last century, and especially with the English and Scotch School of Psychologists, has enabled him to illustrate the subtle speculations of which he treats in a manner calculated to render them more intelligible to the English mind than they can be by writers trained solely in the technicalities of modern German schools. The editor has verified all the references, and added valuable Notes, in which he points out sources of more complete information. The Lectures constitute a History of the Platonic Philosophy—its seed-time, maturity, and decay.

SERMONS AND LETTERS ON ROMANISM.—See THEOLOGICAL CATALOGUE.

Calderwood.—PHILOSOPHY OF THE INFINITE: A Treatise on Man's Knowledge of the Infinite Being, in answer to Sir W. Hamilton and Dr. Mansel. By the Rev. HENRY CALDERWOOD, M.A., LL.D., Professor of Moral Philosophy in the University of Edinburgh. Cheaper Edition. 8vo. 7s. 6d.

The purpose of this volume is, by a careful analysis of consciousness, to prove, in opposition to Sir W. Hamilton and Mr. Mansel, that man possesses a notion of an Infinite Being, and to ascertain the peculiar nature of the conception and the particular relations in which it is found to arise. The province of Faith as related to that of Knowledge, and the characteristics of Knowledge and Thought as bearing on this subject, are examined; and separate chapters are devoted to the consideration of our knowledge of the Infinite as First Cause, as Moral Governor, and as the Object of Worship. "A book of great ability . . . written in a clear style, and may be easily understood by even those who are not versed in such discussions."—British Quarterly Review.

Elam.—A PHYSICIAN'S PROBLEMS. — See MEDICAL CATALOGUE, preceding.

Galton (Francis).—HEREDITARY GENIUS: An Inquiry into its Laws and Consequences. See PHYSICAL SCIENCE CATALOGUE, preceding.

Green (J. H.)—SPIRITUAL PHILOSOPHY: Founded on the Teaching of the late SAMUEL TAYLOR COLERIDGE. By the late JOSEPH HENRY GREEN, F.R.S., D.C.L. Edited, with a Memoir of the Author's Life, by JOHN SIMON, F.R.S., Medical Officer of Her Majesty's Privy Council, and Surgeon to St. Thomas's Hospital. Two Vols. 8vo. 25s.

The late Mr. Green, the eminent surgeon, was for many years the intimate friend and disciple of Coleridge, and an ardent student of philosophy. The language of Coleridge's will imposed on Mr. Green the obligation of devoting, so far as necessary, the remainder of his life to the one task of systematising, developing, and establishing the doctrines of the Coleridgean philosophy. With the assistance of Coleridge's manuscripts, but especially from the knowledge he possessed of Coleridge's doctrines, and independent study of at least the basal principles and metaphysics of the sciences and of all the phenomena of human life, he proceeded logically to work out a system of universal philosophy such as he deemed would in the main accord with his master's aspirations. After many years of preparatory labour he resolved to complete in a compendious form a work which should give in system the doctrines most distinctly Coleridgean. The result is these two volumes. The first volume

is devoted to the general principles of philosophy; the second aims at vindicating à priori (on principles for which the first volume has contended) the essential doctrines of Christianity. The work is divided into four parts: I. "On the Intellectual Faculties and processes which are concerned in the Investigation of Truth." II. "Of First Principles in Philosophy." III. "Truths of Religion." IV. "The Idea of Christianity in relation to Controversial Philosophy."

Huxley (Professor.)—LAY SERMONS, ADDRESSES, AND REVIEWS. See PHYSICAL SCIENCE CATALOGUE, preceding.

Jevons.—Works by W. STANLEY JEVONS, M.A., Professor of Logic in Owens College, Manchester :—

THE SUBSTITUTION OF SIMILARS, the True Principle of Reasoning. Derived from a Modification of Aristotle's Dictum. Fcap. 8vo. 2s. 6d.

"All acts of reasoning," the author says, "seem to me to be different cases of one uniform process, which may perhaps be best described as the substitution of similars. This phrase clearly expresses that familiar mode in which we continually argue by analogy from like to like, and take one thing as a representative of another. The chief difficulty consists in showing that all the forms of the old logic, as well as the fundamental rules of mathematical reasoning, may be explained upon the same principle; and it is to this difficult task I have devoted the most attention. Should my notion be true, a vast mass of technicalities may be swept from our logical text-books and yet the small remaining part of logical doctrine will prove far more useful than all the learning of the Schoolmen." Prefixed is a plan of a new reasoning machine, the Logical Abacus, the construction and working of which is fully explained in the text and Appendix. "Mr. Jevons' book is very clear and intelligible, and quite worth consulting."—Guardian.

ELEMENTARY LESSONS IN LOGIC.—See EDUCATIONAL CATALOGUE.

Maccoll.—THE GREEK SCEPTICS, from Pyrrho to Sextus. An Essay which obtained the Hare Prize in the year 1868. By

NORMAN MACCOLL, B.A., Scholar of Downing College, Cambridge. Crown 8vo, 3s. 6d.

This Essay consists of five parts: I. "Introduction." II. "Pyrrho and Timon." III. "The New Academy." IV. "The Later Sceptics." V. "The Pyrrhoneans and New Academy contrasted."—"Mr. Maccoll has produced a monograph which merits the gratitude of all students of philosophy. His style is clear and vigorous; he has mastered the authorities, and criticises them in a modest but independent spirit."—Pall Mall Gazette.

M'Cosh.—Works by JAMES M'COSH, LL.D., President of Princeton College, New Jersey, U.S.

"He certainly shows himself skilful in that application of logic to psychology, in that inductive science of the human mind which is the fine side of English philosophy. His philosophy as a whole is worthy of attention."—Revue de Deux Mondes.

THE METHOD OF THE DIVINE GOVERNMENT, Physical and Moral. Tenth Edition. 8vo. 10s. 6d.

This work is divided into four books. The first presents a general view of the Divine Government as fitted to throw light on the character of God; the second deals with the method of the Divine Government in the physical world; the third treats of the principles of the human mind through which God governs mankind; and the fourth is on Pastoral and Revealed Religion, and the Restoration of Man. An Appendix, consisting of seven articles, investigates the fundamental principles which underlie the speculations of the treatise. "This work is distinguished from other similar ones by its being based upon a thorough study of physical science, and an accurate knowledge of its present condition, and by its entering in a deeper and more unfettered manner than its predecessors upon the discussion of the appropriate psychological, ethical, and theological questions. The author keeps aloof at once from the à priori idealism and dreaminess of German speculation since Schelling, and from the onesidedness and narrowness of the empiricism and positivism which have so prevailed in England."—Dr. Ulrici, in "Zeitschrift für Philosophie."

THE INTUITIONS OF THE MIND. A New Edition. 8vo. cloth. 10s. 6d.

M'Cosh (J.)—continued.

The object of this treatise is to determine the true nature of Intuition, and to investigate its laws. It starts with a general view of intuitive convictions, their character and the method in which they are employed, and passes on to a more detailed examination of them, treating them under the various heads of "Primitive Cognitions," "Primitive Beliefs," "Primitive Judgments," and "Moral Convictions." Their relations to the various sciences, mental and physical, are then examined. Collateral criticisms are thrown into preliminary and supplementary chapters and sections. "The undertaking to adjust the claims of the sensational and intuitional philosophies, and of the à posteriori and à priori methods, is accomplished in this work with a great amount of success."—Westminster Review. "I value it for its large acquaintance with English Philosophy, which has not led him to neglect the great German works. I admire the moderation and clearness, as well as comprehensiveness, of the author's views."—Dr. Dorner, of Berlin.

AN EXAMINATION OF MR. J. S. MILL'S PHILOSOPHY:
Being a Defence of Fundamental Truth. Crown 8vo. 7s. 6d.

This volume is not put forth by its author as a special reply to Mr. Mill's "Examination of Sir William Hamilton's Philosophy." In that work Mr. Mill has furnished the means of thoroughly estimating his theory of mind, of which he had only given hints and glimpses in his logical treatise. It is this theory which Dr. M'Cosh professes to examine in this volume; his aim is simply to defend a portion of primary truth which has been assailed by an acute thinker who has extensive influence in England. "In such points as Mr. Mill's notions of intuitions and necessity, he will have the voice of mankind with him."—Athenæum. "Such a work greatly needed to be done, and the author was the man to do it. This volume is important, not merely in reference to the views of Mr. Mill, but of the whole school of writers, past and present, British and Continental, he so ably represents."—Princeton Review.

THE LAWS OF DISCURSIVE THOUGHT: Being a Text-
book of Formal Logic. Crown 8vo. 5s.

The main feature of this Logical Treatise is to be found in the more thorough investigation of the nature of the notion, in regard to

M'Cosh (J.)—*continued.*

which the views of the school of Locke and Whately are regarded by the author as very defective, and the views of the school of Kant and Hamilton altogether erroneous. The author believes that errors spring far more frequently from obscure, inadequate, indistinct, and confused Notions, and from not placing the Notions in their proper relation in judgment, than from Ratiocination. In this treatise, therefore, the Notion (with the term, and the Relation of Thought to Language) will be found to occupy a larger relative place than in any logical work written since the time of the famous Art of Thinking. "The amount of summarized information which it contains is very great; and it is the only work on the very important subject with which it deals. Never was such a work so much needed as in the present day."—London Quarterly Review.

CHRISTIANITY AND POSITIVISM : A Series of Lectures to the Times on Natural Theology and Apologetics. Crown 8vo. 7s. 6d.

These Lectures were delivered in New York, by appointment, in the beginning of 1871, as the second course on the foundation of the Union Theological Seminary. There are ten Lectures in all, divided into three series :—I. "Christianity and Physical Science" (three lectures). II. "Christianity and Mental Science" (four lectures). III. "Christianity and Historical Investigation" (three lectures). The Appendix contains articles on "Gaps in the Theory of Development;" "Darwin's Descent of Man." "Principles of Herbert Spencer's Philosophy." In the course of the Lectures Dr. M'Cosh discusses all the most important scientific problems which are supposed to affect Christianity.

Masson.—RECENT BRITISH PHILOSOPHY : A Review, with Criticisms ; including some Comments on Mr. Mill's Answer to Sir William Hamilton. By DAVID MASSON, M.A., Professor of Rhetoric and English Literature in the University of Edinburgh. Crown 8vo. 6s.

The author, in his usual graphic and forcible manner, reviews in considerable detail, and points out the drifts of the philosophical speculations of the previous thirty years, bringing under notice the work of all the principal philosophers who have been at work during

Masson (D.)—*continued.*

that period on the highest problems which concern humanity. The four chapters are thus titled:—I. "A Survey of Thirty Years." II. "The Traditional Differences: how repeated in Carlyle, Hamilton, and Mill." III. "Effects of Recent Scientific Conceptions on Philosophy." IV. "Latest Drifts and Groupings." The last seventy-six pages are devoted to a Review of Mr. Mill's criticism of Sir William Hamilton's Philosophy. "We can nowhere point to a work which gives so clear an exposition of the course of philosophical speculation in Britain during the past century, or which indicates so instructively the mutual influences of philosophic and scientific thought."—Fortnightly Review.

BRITISH NOVELISTS.—See BEILLES LETTRES CATALOGUE.

LIFE OF MILTON.—See BIOGRAPHICAL CATALOGUE.

Maudsley.—Works by HENRY MAUDSLEY, M.D., Professor of Medical Jurisprudence in University College, London:—

BODY AND MIND: An Inquiry into their Connection and Mutual Influence, specially in reference to Mental Diseases. See MEDICAL CATALOGUE, preceding.

THE PHYSIOLOGY AND PATHOLOGY OF MIND.
See MEDICAL CATALOGUE, preceding.

Maurice.—Works by the Rev. FREDERICK DENISON MAURICE, M.A., Professor of Moral Philosophy in the University of Cambridge. (For other Works by the same Author, see THEOLOGICAL CATALOGUE.)

SOCIAL MORALITY. Twenty-one Lectures delivered in the University of Cambridge. 8vo. 14s.

In this series of Lectures, Professor Maurice considers, historically and critically, Social Morality in its three main aspects: I. "The Relations which spring from the Family—Domestic Morality." II. "The Relations which subsist among the various constituents of a Nation—National Morality." III. "As it concerns Universal Humanity—Universal Morality." Appended to each series is a chapter on "Worship:" first, "Family Worship;" second,

Maurice (F. D.)—*continued.*

“National Worship;” third, “Universal Worship.” “Whilst reading it we are charmed by the freedom from exclusiveness and prejudice, the large charity, the loftiness of thought, the eagerness to recognize and appreciate whatever there is of real worth extant in the world, which animates it from one end to the other. We gain new thoughts and new ways of viewing things, even more, perhaps, from being brought for a time under the influence of so noble and spiritual a mind.”—Athenæum.

THE CONSCIENCE: Lectures on Casuistry, delivered in the University of Cambridge. New and Cheaper Edition. Crown 8vo. 5s.

In this series of nine Lectures, Professor Maurice, with his wonted force and breadth and freshness, endeavours to settle what is meant by the word “Conscience,” and discusses the most important questions immediately connected with the subject. Taking “Casuistry” in its old sense as being the “study of cases of Conscience,” he endeavours to show in what way it may be brought to bear at the present day upon the acts and thoughts of our ordinary existence. He shows that Conscience asks for laws, not rules; for freedom, not chains; for education, not suppression. He has abstained from the use of philosophical terms, and has touched on philosophical systems only when he fancied “they were interfering with the rights and duties of wayfarers.” The Saturday Review says: “We rise from them with detestation of all that is selfish and mean, and with a living impression that there is such a thing as goodness after all.”

MORAL AND METAPHYSICAL PHILOSOPHY. New Edition and Preface. Vol. I. Ancient Philosophy and the First to the Thirteenth Centuries; Vol. II. the Fourteenth Century and the French Revolution, with a glimpse into the Nineteenth Century. 2 Vols. 8vo. 25s.

This is an Edition in two volumes of Professor Maurice’s History of Philosophy from the earliest period to the present time. It was formerly scattered throughout a number of separate volumes, and it is believed that all admirers of the author and all students of philosophy will welcome this compact Edition. The subject is one of the highest importance, and it is treated here with fulness and

candour, and in a clear and interesting manner. In a long introduction to this Edition, in the form of a dialogue, Professor Maurice justifies some of his own peculiar views, and touches upon some of the most important topics of the time.

Murphy.—HABIT AND INTELLIGENCE, in Connection with the Laws of Matter and Force : A Series of Scientific Essays. By JOSEPH JOHN MURPHY. Two Vols. 8vo. 16s.

The author's chief purpose in this work has been to state and to discuss what he regards as the special and characteristic principles of life. The most important part of the work treats of those vital principles which belong to the inner domain of life itself, as distinguished from the principles which belong to the border-land where life comes into contact with inorganic matter and force. In the inner domain of life we find two principles, which are, the author believes, coextensive with life and peculiar to it : these are Habit and Intelligence. He has made as full a statement as possible of the laws under which habits form, disappear, alter under altered circumstances, and vary spontaneously. He discusses that most important of all questions, whether intelligence is an ultimate fact, incapable of being resolved into any other, or only a resultant from the laws of habit. The latter part of the first volume is occupied with the discussion of the question of the Origin of Species. The first part of the second volume is occupied with an inquiry into the process of mental growth and development, and the nature of mental intelligence. In the chapter that follows, the author discusses the science of history, and the three concluding chapters contain some ideas on the classification, the history, and the logic, of the sciences. The author's aim has been to make the subjects treated of intelligible to any ordinary intelligent man. "We are pleased to listen," says the Saturday Review, "to a writer who has so firm a foothold upon the ground within the scope of his immediate survey, and who can enunciate with so much clearness and force propositions which come within his grasp."

Thring (E., M.A.)—THOUGHTS ON LIFE-SCIENCE.

By EDWARD THRING, M.A. (Benjamin Place), Head Master of Uppingham School. New Edition, enlarged and revised. Crown 8vo. 7s. 6d.

In this volume are discussed in a familiar manner some of the most interesting problems between Science and Religion, Reason and

Feeling. "Learning and Science," says the author, "are claiming the right of building up and pulling down everything, especially the latter. It has seemed to me no useless task to look steadily at what has happened, to take stock as it were of men's gains, and to endeavour amidst new circumstances to arrive at some rational estimate of the bearings of things, so that the limits of what is possible at all events may be clearly marked out for ordinary readers. . . . This book is an endeavour to bring out some of the main facts of the world."

Venn.—THE LOGIC OF CHANCE: An Essay on the Foundations and Province of the Theory of Probability, with especial reference to its application to Moral and Social Science. By JOHN VENN, M.A., Fellow of Gonville and Caius College, Cambridge. Fcap. 8vo. 7s. 6d.

This Essay is in no sense mathematical. Probability, the author thinks, may be considered to be a portion of the province of Logic regarded from the material point of view. The principal objects of this Essay are to ascertain how great a portion it comprises, where we are to draw the boundary between it and the contiguous branches of the general science of evidence, what are the ultimate foundations upon which its rules rest, what the nature of the evidence they are capable of affording, and to what class of subjects they may most fitly be applied. The general design of the Essay, as a special treatise on Probability, is quite original, the author believing that erroneous notions as to the real nature of the subject are disastrously prevalent. "Exceedingly well thought and well written," says the Westminster Review. The Nonconformist calls it a "masterly book."

